

Terminal Transferase



1-800-632-7799
info@neb.com
www.neb.com



M0315S 021160618061

M0315S



500 units **20,000 U/ml** **Lot: 0211606**
RECOMBINANT **Store at -20°C** **Exp: 6/18**

Description: Terminal Transferase (TdT) is a template independent polymerase that catalyzes the addition of deoxynucleotides to the 3' hydroxyl terminus of DNA molecules. Protruding, recessed or blunt-ended double or single-stranded DNA molecules serve as a substrate for TdT. The 58.3 kDa enzyme does not have 5' or 3' exonuclease activity. The addition of Co²⁺ in the reaction makes tailing more efficient.

New Reaction Buffer

Terminal Transferase



1-800-632-7799
info@neb.com
www.neb.com



M0315S 021160618061

M0315S



500 units **20,000 U/ml** **Lot: 0211606**
RECOMBINANT **Store at -20°C** **Exp: 6/18**

Description: Terminal Transferase (TdT) is a template independent polymerase that catalyzes the addition of deoxynucleotides to the 3' hydroxyl terminus of DNA molecules. Protruding, recessed or blunt-ended double or single-stranded DNA molecules serve as a substrate for TdT. The 58.3 kDa enzyme does not have 5' or 3' exonuclease activity. The addition of Co²⁺ in the reaction makes tailing more efficient.

New Reaction Buffer

Source: An *E. coli* strain that carries the cloned Terminal Transferase gene from calf thymus.

Applications:

- Addition of homopolymer tails to the 3' ends of DNA
- Labeling the 3' ends of DNA with modified nucleotides (e.g., ddNTP, DIG-dUTP)
- TUNEL assay (*in situ* localization of apoptosis)
- TdT dependent PCR

Supplied in: 50 mM KPO₄ (pH 7.3 @ 25°C), 100 mM NaCl, 1.43 mM 2-mercaptoethanol, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:

10X Terminal Transferase Reaction Buffer, 10X (2.5 mM) solution of CoCl₂.

Reaction Conditions: 1X Terminal Transferase Reaction Buffer, supplemented with 0.25 mM CoCl₂. Incubate at 37°C.

1X Terminal Transferase Reaction Buffer:

50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
pH 7.9 @ 25°C

Source: An *E. coli* strain that carries the cloned Terminal Transferase gene from calf thymus.

Applications:

- Addition of homopolymer tails to the 3' ends of DNA
- Labeling the 3' ends of DNA with modified nucleotides (e.g., ddNTP, DIG-dUTP)
- TUNEL assay (*in situ* localization of apoptosis)
- TdT dependent PCR

Supplied in: 50 mM KPO₄ (pH 7.3 @ 25°C), 100 mM NaCl, 1.43 mM 2-mercaptoethanol, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:

10X Terminal Transferase Reaction Buffer, 10X (2.5 mM) solution of CoCl₂.

Reaction Conditions: 1X Terminal Transferase Reaction Buffer, supplemented with 0.25 mM CoCl₂. Incubate at 37°C.

1X Terminal Transferase Reaction Buffer:

50 mM potassium acetate
20 mM Tris-acetate
10 mM magnesium acetate
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme catalyzing the incorporation of 1 nmol dATP into acid-insoluble material in a total reaction volume of 1 ml in 1 hour at 37°C using d(A)₁₈ as a primer.

Unit Assay Conditions: 1X Terminal Transferase Reaction Buffer, 0.72 μM d(A)₁₈, 0.2 mM dATP and 1.0 μCi [³H]-dATP in a 50 μl total reaction volume.

Quality Control Assays

Exonuclease Activity: Incubation of 50 units of enzyme with 1 μg sonicated [³H] DNA (2 × 10⁵ cpm/μg) for 4 hours at 37°C in 50 μl assay buffer released < 0.5% radioactivity.

Endonuclease Activity: Incubation of 50 units of enzyme with 1 μg φX174 RF I DNA for 4 hours at 37°C in a 50 μl reaction buffer resulted in < 10% conversion to RF II.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Heat Inactivation: 75°C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme catalyzing the incorporation of 1 nmol dATP into acid-insoluble material in a total reaction volume of 1 ml in 1 hour at 37°C using d(A)₁₈ as a primer.

Unit Assay Conditions: 1X Terminal Transferase Reaction Buffer, 0.72 μM d(A)₁₈, 0.2 mM dATP and 1.0 μCi [³H]-dATP in a 50 μl total reaction volume.

Quality Control Assays

Exonuclease Activity: Incubation of 50 units of enzyme with 1 μg sonicated [³H] DNA (2 × 10⁵ cpm/μg) for 4 hours at 37°C in 50 μl assay buffer released < 0.5% radioactivity.

Endonuclease Activity: Incubation of 50 units of enzyme with 1 μg φX174 RF I DNA for 4 hours at 37°C in a 50 μl reaction buffer resulted in < 10% conversion to RF II.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Heat Inactivation: 75°C for 20 minutes.

A Typical DNA Tailing Reaction:

1. Mix:
 - a. 5.0 μl 10X TdT Buffer
 - b. 5.0 μl 2.5 mM CoCl₂ solution provided
 - c. 5.0 pmols DNA (330 ng for 100 bp, 1 μg for 300 bp, 10 pmols DNA ends)*
 - d. 0.5 μl 10 mM dNTP (alpha-³²P dATP may also be used)
 - e. 0.5 μl Terminal Transferase (20 units/μl) deionized H₂O to a final volume of 50 μl.
2. Incubate at 37°C for 30 minutes.
3. Stop the reaction by heating to 70°C for 10 minutes or by adding 10 μl of 0.2 M EDTA (pH 8.0).

*To determine approximate amount of DNA (ng/pmol), multiply the number of base pairs by 0.66. Example: 300 bp × 0.66 = 198 ng/pmol. For 5.0 pmols multiply by 5, resulting in 990 ng/5 pmol.

The table on the reverse side can be used as a guide (values are approximate and are given for a 30 minutes incubation at 37°C in the recommended buffer).

(see other side)

CERTIFICATE OF ANALYSIS

A Typical DNA Tailing Reaction:

1. Mix:
 - a. 5.0 μl 10X TdT Buffer
 - b. 5.0 μl 2.5 mM CoCl₂ solution provided
 - c. 5.0 pmols DNA (330 ng for 100 bp, 1 μg for 300 bp, 10 pmols DNA ends)*
 - d. 0.5 μl 10 mM dNTP (alpha-³²P dATP may also be used)
 - e. 0.5 μl Terminal Transferase (20 units/μl) deionized H₂O to a final volume of 50 μl.
2. Incubate at 37°C for 30 minutes.
3. Stop the reaction by heating to 70°C for 10 minutes or by adding 10 μl of 0.2 M EDTA (pH 8.0).

*To determine approximate amount of DNA (ng/pmol), multiply the number of base pairs by 0.66. Example: 300 bp × 0.66 = 198 ng/pmol. For 5.0 pmols multiply by 5, resulting in 990 ng/5 pmol.

The table on the reverse side can be used as a guide (values are approximate and are given for a 30 minutes incubation at 37°C in the recommended buffer).

(see other side)

CERTIFICATE OF ANALYSIS

The rate of addition of dNTP's and thus the length of the tail is a function of the ratio of 3' DNA ends: dNTP concentration, and also which dNTP is used.

DNA Tailing Guide:

pmols 3' ends pmol dNTP	Tail Length			
	dA	dC	dG	dT
1:100	1–5	1–3	1–3	1–5
1:1,000	10–20	10–20	5–10	10–20
1:5,000	100–300	50–200	10–25	200–300

References:

1. Chang, L.M. and Bollum, F.J. (1986) *CRC Crit. Rev. Biochem.* 21, 27–52.
2. Roychoudhury, R., Jay, E. and Wu R. (1976) *Nucl. Acids Res.* 3, 101–116.
3. Tu, C.-P.D. and Cohen, S.N. (1980) *Gene* 10, 177–183.
4. Boule, J.B., Rougeon, F. and Papanicolaou C. (2001) *J. Biol. Chem.* 276, 31388–31393.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

The rate of addition of dNTP's and thus the length of the tail is a function of the ratio of 3' DNA ends: dNTP concentration, and also which dNTP is used.

DNA Tailing Guide:

pmols 3' ends pmol dNTP	Tail Length			
	dA	dC	dG	dT
1:100	1–5	1–3	1–3	1–5
1:1,000	10–20	10–20	5–10	10–20
1:5,000	100–300	50–200	10–25	200–300

References:

1. Chang, L.M. and Bollum, F.J. (1986) *CRC Crit. Rev. Biochem.* 21, 27–52.
2. Roychoudhury, R., Jay, E. and Wu R. (1976) *Nucl. Acids Res.* 3, 101–116.
3. Tu, C.-P.D. and Cohen, S.N. (1980) *Gene* 10, 177–183.
4. Boule, J.B., Rougeon, F. and Papanicolaou C. (2001) *J. Biol. Chem.* 276, 31388–31393.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.