

hSMUG1



1-800-632-7799
info@neb.com
www.neb.com



M0336S 001160517051

M0336S

500 units 5,000 U/ml Lot: 0011605

RECOMBINANT Store at -20° Exp: 5/17

Description: Human single-strand-selective monofunctional uracil-DNA Glycosylase SMUG1 excises deoxyuracil and deoxyuracil-derivatives bearing an oxidized group at C5, such as 5-hydroxyuracil, 5-hydroxymethyluracil and 5-formyluracil in ssDNA and dsDNA (1,2,3).

Source: An *E. coli* strain which carries the cloned human SMUG1 gene.

Applications:

- Oxidative DNA damage studies
- Single cell gel electrophoresis (comet assay) (4,5,6)

hSMUG1



1-800-632-7799
info@neb.com
www.neb.com



M0336S 001160517051

M0336S

500 units 5,000 U/ml Lot: 0011605

RECOMBINANT Store at -20° Exp: 5/17

Description: Human single-strand-selective monofunctional uracil-DNA Glycosylase SMUG1 excises deoxyuracil and deoxyuracil-derivatives bearing an oxidized group at C5, such as 5-hydroxyuracil, 5-hydroxymethyluracil and 5-formyluracil in ssDNA and dsDNA (1,2,3).

Source: An *E. coli* strain which carries the cloned human SMUG1 gene.

Applications:

- Oxidative DNA damage studies
- Single cell gel electrophoresis (comet assay) (4,5,6)

Supplied in: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 1, 100X BSA.

Reaction Conditions: 1X NEBuffer 1 supplemented with 100 µg/ml BSA. Incubate at 37°C.

1X NEBuffer 1:

10 mM Bis Tris Propane-HCl
10 mM MgCl₂
1 mM DTT
pH 7.0 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to excise 1 pmol of deoxyuracil from a 34 mer oligonucleotide duplex containing a single dU site in a total reaction volume of 10 µl in 1 hour at 37°C.

Molecular Weight: 29,861 Daltons

Diluent Compatibility: Diluent C

250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol.

Supplied in: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X NEBuffer 1, 100X BSA.

Reaction Conditions: 1X NEBuffer 1 supplemented with 100 µg/ml BSA. Incubate at 37°C.

1X NEBuffer 1:

10 mM Bis Tris Propane-HCl
10 mM MgCl₂
1 mM DTT
pH 7.0 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to excise 1 pmol of deoxyuracil from a 34 mer oligonucleotide duplex containing a single dU site in a total reaction volume of 10 µl in 1 hour at 37°C.

Molecular Weight: 29,861 Daltons

Diluent Compatibility: Diluent C

250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 0.15% Triton X-100, 200 µg/ml BSA and 50% glycerol.

Unit Assay Conditions: 1X NEBuffer 1 supplemented with 100 µg/ml BSA containing 5 pmol of fluorescently labeled oligonucleotide duplex in a total reaction volume of 10 µl. After release of dU, the oligonucleotide is cleaved by treating the resulting AP site with 100 mM NaOH for 10 minutes at 80°C.

Quality Control Assays

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 50 units of hSMUG1 incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 20 µl reaction containing 50 units of hSMUG1 with a 10 nM mixture of single-stranded and double-stranded fluorescently labeled oligonucleotides containing blunt ends, 5' extensions, and 3' extensions for 30 minutes at 37°C yields no detectable degradation as determined by high resolution capillary electrophoresis.

Unit Assay Conditions: 1X NEBuffer 1 supplemented with 100 µg/ml BSA containing 5 pmol of fluorescently labeled oligonucleotide duplex in a total reaction volume of 10 µl. After release of dU, the oligonucleotide is cleaved by treating the resulting AP site with 100 mM NaOH for 10 minutes at 80°C.

Quality Control Assays

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 50 units of hSMUG1 incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 20 µl reaction containing 50 units of hSMUG1 with a 10 nM mixture of single-stranded and double-stranded fluorescently labeled oligonucleotides containing blunt ends, 5' extensions, and 3' extensions for 30 minutes at 37°C yields no detectable degradation as determined by high resolution capillary electrophoresis.

Endonuclease Activity: Incubation of a 50 µl reaction containing 50 units of hSMUG1 with 1 µg φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

Heat Inactivation: 20 minutes at 65°C.

Usage Note: hSMUG1 has 50% activity on 5-hydroxymethyluracil when compared to uracil. hSMUG1 has 50% activity on ssDNA compared to ds DNA.

References:

1. Cannon-Crlson, S.V. et al. (1989) *J. Biol. Chem.* 264, 13306-13312.
2. Masaoka, A. et al. (2003) *Biochemistry* 42, 5003-5012.
3. Wibley, J.E.A. et al. (2003) *Cell* 11, 1647-1659.
4. Singh, N. et al. (1988) *Experimental Cell Research* 175, 184-191.
5. Collins, A. et al. (1993) *Carcinogenesis* 14, 1733-1735.
6. Collins, A. et al. (1996) *Environmental Health Perspectives* 104, 465-469.

CERTIFICATE OF ANALYSIS

Endonuclease Activity: Incubation of a 50 µl reaction containing 50 units of hSMUG1 with 1 µg φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

Heat Inactivation: 20 minutes at 65°C.

Usage Note: hSMUG1 has 50% activity on 5-hydroxymethyluracil when compared to uracil. hSMUG1 has 50% activity on ssDNA compared to ds DNA.

References:

1. Cannon-Crlson, S.V. et al. (1989) *J. Biol. Chem.* 264, 13306-13312.
2. Masaoka, A. et al. (2003) *Biochemistry* 42, 5003-5012.
3. Wibley, J.E.A. et al. (2003) *Cell* 11, 1647-1659.
4. Singh, N. et al. (1988) *Experimental Cell Research* 175, 184-191.
5. Collins, A. et al. (1993) *Carcinogenesis* 14, 1733-1735.
6. Collins, A. et al. (1996) *Environmental Health Perspectives* 104, 465-469.

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.