

Exonuclease VII



M0379S

200 units **10,000 units/ml** **Lot: 0021703**
RECOMBINANT **Store at -20°C** **Exp: 3/19**

Description: Exonuclease VII, (Exo VII) derived from *E. coli*, cleaves single-stranded DNA (ssDNA) from both 5'→3' and 3'→5' direction. This enzyme is not active on linear or circular dsDNA (1,2). It is useful for removal of single stranded oligonucleotide primers (3) from a completed PCR reaction when different primers are required for subsequent PCR reactions. Digestion of ssDNA by Exonuclease VII is metal-independent.

Source: An *E. coli* strain that carries cloned Exonuclease VII (XseA and XseB) genes from *E. coli*.

Exonuclease VII



M0379S

200 units **10,000 units/ml** **Lot: 0021703**
RECOMBINANT **Store at -20°C** **Exp: 3/19**

Description: Exonuclease VII, (Exo VII) derived from *E. coli*, cleaves single-stranded DNA (ssDNA) from both 5'→3' and 3'→5' direction. This enzyme is not active on linear or circular dsDNA (1,2). It is useful for removal of single stranded oligonucleotide primers (3) from a completed PCR reaction when different primers are required for subsequent PCR reactions. Digestion of ssDNA by Exonuclease VII is metal-independent.

Source: An *E. coli* strain that carries cloned Exonuclease VII (XseA and XseB) genes from *E. coli*.

Applications:

- Removal of single-stranded oligonucleotide primers after PCR (3)
- Removal of terminal phosphorothioated ss-oligonucleotide primers after PCR
- Mapping positions of introns in genomic DNA (4)
- Removal of single-stranded DNA from dsDNA

Supplied in: 50% glycerol containing 50 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM DTT, 0.1 M NaCl and 0.1% Triton™ X-100. (pH 7.5 @ 25°C).

Reagents Supplied with Enzyme:
5X Exonuclease VII Reaction Buffer.

Reaction Conditions: 1X Exonuclease VII Reaction Buffer. Incubate at 37°C.

1X Exonuclease VII Reaction Buffer:
50 mM Tris-HCl
50 mM sodium phosphate
10 mM 2-mercaptoethanol
8 mM EDTA
(pH 8.0 @ 25°C)

Applications:

- Removal of single-stranded oligonucleotide primers after PCR (3)
- Removal of terminal phosphorothioated ss-oligonucleotide primers after PCR
- Mapping positions of introns in genomic DNA (4)
- Removal of single-stranded DNA from dsDNA

Supplied in: 50% glycerol containing 50 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM DTT, 0.1 M NaCl and 0.1% Triton™ X-100. (pH 7.5 @ 25°C).

Reagents Supplied with Enzyme:
5X Exonuclease VII Reaction Buffer.

Reaction Conditions: 1X Exonuclease VII Reaction Buffer. Incubate at 37°C.

1X Exonuclease VII Reaction Buffer:
50 mM Tris-HCl
50 mM sodium phosphate
10 mM 2-mercaptoethanol
8 mM EDTA
(pH 8.0 @ 25°C)

Unit Definition: One unit is defined as the amount of enzyme that will catalyze the release of 1 nmol of acid-soluble nucleotide in a total reaction volume of 50 µl in 30 minutes at 37°C.

Unit Assay Conditions: 1X Exonuclease VII Reaction Buffer containing 0.15 mM sonicated single-stranded [³H] *E. coli* DNA.

Quality Control Assays

16-Hour Incubation: A 50 µl reaction in NEBuffer 4 containing 1 µg of HaeIII-cut φX174 Phage DNA and 10 units of Exonuclease VII incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

ds-Exonuclease Activity: Incubation of a 50 µl reaction containing 10 units of Exonuclease VII in NEBuffer 4 with 1 µg of double-stranded [³H] *E. coli* DNA (10⁵ cpm/µg) for 4 hours at 37°C released < 0.5% of the total radioactivity.

Unit Definition: One unit is defined as the amount of enzyme that will catalyze the release of 1 nmol of acid-soluble nucleotide in a total reaction volume of 50 µl in 30 minutes at 37°C.

Unit Assay Conditions: 1X Exonuclease VII Reaction Buffer containing 0.15 mM sonicated single-stranded [³H] *E. coli* DNA.

Quality Control Assays

16-Hour Incubation: A 50 µl reaction in NEBuffer 4 containing 1 µg of HaeIII-cut φX174 Phage DNA and 10 units of Exonuclease VII incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

ds-Exonuclease Activity: Incubation of a 50 µl reaction containing 10 units of Exonuclease VII in NEBuffer 4 with 1 µg of double-stranded [³H] *E. coli* DNA (10⁵ cpm/µg) for 4 hours at 37°C released < 0.5% of the total radioactivity.

ds-Endonuclease Activity: Incubation of 50 units of Exonuclease VII with 1 µg φX174 RF I DNA for 4 hours at 37°C in NEBuffer 4 resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

ss-Endonuclease Activity: Incubation of 10 units of Exonuclease VII with 1 µg of M13 ssDNA for 1 hour at 37°C in NEBuffer 4 resulted in < 20% decrease in the amount of closed circular DNA as determined by agarose gel electrophoresis.

RNase Activity (Extended Digestion): A 10 µl reaction in NEBuffer 4 containing 40 ng of F-300 RNA probe and a minimum of 10 unit of Exonuclease VII is incubated at 37°C. After incubation for 4 hours, > 90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescein imaging detection

Physical Purity: Purified to > 99% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

(see other side)

CERTIFICATE OF ANALYSIS

ds-Endonuclease Activity: Incubation of 50 units of Exonuclease VII with 1 µg φX174 RF I DNA for 4 hours at 37°C in NEBuffer 4 resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

ss-Endonuclease Activity: Incubation of 10 units of Exonuclease VII with 1 µg of M13 ssDNA for 1 hour at 37°C in NEBuffer 4 resulted in < 20% decrease in the amount of closed circular DNA as determined by agarose gel electrophoresis.

RNase Activity (Extended Digestion): A 10 µl reaction in NEBuffer 4 containing 40 ng of F-300 RNA probe and a minimum of 10 unit of Exonuclease VII is incubated at 37°C. After incubation for 4 hours, > 90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescein imaging detection

Physical Purity: Purified to > 99% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

(see other side)

CERTIFICATE OF ANALYSIS

qPCR DNA Contamination (*E. coli* Genomic):

A minimum of 10 units of Exonuclease VII is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is less than 1 copy of *E. coli* genome.

Heat Inactivation: 95°C for 10 minutes.

References:

1. Chase, et al. (1974) *J. Biol. Chem.* 249, 4545–4552.
2. Chase, et al. (1974) *J. Biol. Chem.* 249, 4553–4561.
3. Li, H. et al. (1991) *Nucl. Acids Res.* 19, 3139–3141.
4. Berk, A.J. et al. (1978) *Proc. Natl. Acad. Sci. USA*, 75, 1274–1278.

Page 2 (M0379)



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

SYBR® is a registered trademark of Life Technologies, Inc.

TRITON™ is a trademark of The Dow Chemical Company or an affiliated company of Dow.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

qPCR DNA Contamination (*E. coli* Genomic):

A minimum of 10 units of Exonuclease VII is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is less than 1 copy of *E. coli* genome.

Heat Inactivation: 95°C for 10 minutes.

References:

1. Chase, et al. (1974) *J. Biol. Chem.* 249, 4545–4552.
2. Chase, et al. (1974) *J. Biol. Chem.* 249, 4553–4561.
3. Li, H. et al. (1991) *Nucl. Acids Res.* 19, 3139–3141.
4. Berk, A.J. et al. (1978) *Proc. Natl. Acad. Sci. USA*, 75, 1274–1278.

Page 2 (M0379)



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

SYBR® is a registered trademark of Life Technologies, Inc.

TRITON™ is a trademark of The Dow Chemical Company or an affiliated company of Dow.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.