

ET SSB



1-800-632-7799
info@neb.com
www.neb.com



M2401S 003150917091

M2401S



50 µg 500 µg/ml Lot: 0031509

RECOMBINANT Store at -20°C Exp: 9/17

Description: ET SSB (Extreme Thermostable Single-Stranded DNA Binding Protein) is a single-stranded DNA binding protein isolated from a hyperthermophilic microorganism, which remains fully active after incubation at 95°C for 60 minutes. Due to the extreme thermostability, ET SSB can be used in applications that require extremely high temperature conditions, such as nucleic acid amplification and sequencing.

Source: An *E. coli* strain that carries the cloned *ssb* gene from a hyperthermophilic organism.

Applications:

- Improve the processivity of DNA polymerase (1)
- Stabilization and marking of ssDNA structure (2)
- Increase the yield and specificity of PCR reactions (3–7)
- Increase the yield and processivity of RT during RT-PCR (8–9)
- Improve DNA sequencing through regions with strong secondary structure (6)
- Enhance the RecA activity for ssDNA binding and strand transfer (10,11)

Supplied in: 20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 0.5 mM DTT, 1 mM EDTA and 50% glycerol.

Unit Definition: Sold by mass of pure protein as determined by OD_{280} ($A_{280} = 0.774$ at 1 mg/ml, 1cm).

Molecular Weight: 16 kDa.

Quality Assurance: ET SSB is purified free of contaminating endonucleases and exonucleases. Each lot is tested for ssDNA binding activity and is visually determined to be > 95% pure on an SDS-polyacrylamide gel.

Quality Control Assays

Exonuclease Activity: Incubation of 20 µg ET SSB for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate, 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of 5 µg ET SSB for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate, 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg φX174 RF I DNA gave < 20% conversion to RF II.

Nuclease Activity: Incubation of 2.5 µg ET SSB for 16 hours at 65°C in 50 µl of reaction buffer containing 50 mM potassium acetate, 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg λ DNA yielded a clear and sharp band on an agarose gel.

Notes On Use: ET SSB is active in any polymerase buffer. Add 200 ng of ET SSB per 50 µl reaction.

References:

1. Myers, T.W. and Romano, L.J. (1988) *J. Biol. Chem.* 263, 17006–17015.
2. Reddy, M.S. (2000) *Biochemistry*, 39, 14250–14262.
3. West, S.C., Cassuto, E. and Howard-Flanders, P. (1982) *Mol. Gen. Genet.* 186, 333–338.
4. Goldmeyer, J., Kong, H. and Tang, W. (2007) *J. Mol. Diagnostics*, 9, 639–644.
5. Delius, H., Mantell, N.J. and Alberts, B. (1972) *J. Mol. Biol.* 67, 341–350.
6. Schwarz, K., Hansen-Hagge, T. and Bartram, C. (1990) *Nucl. Acids Res.* 18, 1079.

(see other side)

CERTIFICATE OF ANALYSIS

ET SSB



1-800-632-7799
info@neb.com
www.neb.com



M2401S 003150917091

M2401S



50 µg 500 µg/ml Lot: 0031509

RECOMBINANT Store at -20°C Exp: 9/17

Description: ET SSB (Extreme Thermostable Single-Stranded DNA Binding Protein) is a single-stranded DNA binding protein isolated from a hyperthermophilic microorganism, which remains fully active after incubation at 95°C for 60 minutes. Due to the extreme thermostability, ET SSB can be used in applications that require extremely high temperature conditions, such as nucleic acid amplification and sequencing.

Source: An *E. coli* strain that carries the cloned *ssb* gene from a hyperthermophilic organism.

Applications:

- Improve the processivity of DNA polymerase (1)
- Stabilization and marking of ssDNA structure (2)
- Increase the yield and specificity of PCR reactions (3–7)
- Increase the yield and processivity of RT during RT-PCR (8–9)
- Improve DNA sequencing through regions with strong secondary structure (6)
- Enhance the RecA activity for ssDNA binding and strand transfer (10,11)

Supplied in: 20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 0.5 mM DTT, 1 mM EDTA and 50% glycerol.

Unit Definition: Sold by mass of pure protein as determined by OD_{280} ($A_{280} = 0.774$ at 1 mg/ml, 1cm).

Molecular Weight: 16 kDa.

Quality Assurance: ET SSB is purified free of contaminating endonucleases and exonucleases. Each lot is tested for ssDNA binding activity and is visually determined to be > 95% pure on an SDS-polyacrylamide gel.

Quality Control Assays

Exonuclease Activity: Incubation of 20 µg ET SSB for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate, 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of 5 µg ET SSB for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate, 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg φX174 RF I DNA gave < 20% conversion to RF II.

Nuclease Activity: Incubation of 2.5 µg ET SSB for 16 hours at 65°C in 50 µl of reaction buffer containing 50 mM potassium acetate, 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg λ DNA yielded a clear and sharp band on an agarose gel.

Notes On Use: ET SSB is active in any polymerase buffer. Add 200 ng of ET SSB per 50 µl reaction.

References:

1. Myers, T.W. and Romano, L.J. (1988) *J. Biol. Chem.* 263, 17006–17015.
2. Reddy, M.S. (2000) *Biochemistry*, 39, 14250–14262.
3. West, S.C., Cassuto, E. and Howard-Flanders, P. (1982) *Mol. Gen. Genet.* 186, 333–338.
4. Goldmeyer, J., Kong, H. and Tang, W. (2007) *J. Mol. Diagnostics*, 9, 639–644.
5. Delius, H., Mantell, N.J. and Alberts, B. (1972) *J. Mol. Biol.* 67, 341–350.
6. Schwarz, K., Hansen-Hagge, T. and Bartram, C. (1990) *Nucl. Acids Res.* 18, 1079.

(see other side)

CERTIFICATE OF ANALYSIS

7. Chou, Q. (1992) *Nucl. Acids Res.* 20, 4371.
8. Oshima, R.G. (1992) *Biotechniques*, 13, 188.
9. Rapley, R. (1994) *Mol. Biotechnol.* 2, 295–298.
10. Olszewski, M. et al. (2005) *Mol. Cell Probes*, 19, 203–205.
11. Baugh, L.R. et al. (2001) *Nucl. Acids Res.* 29, e29.
12. Villalva, C. et al. (2001) *Biotechniques* 31, 81–83, 86.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

7. Chou, Q. (1992) *Nucl. Acids Res.* 20, 4371.
8. Oshima, R.G. (1992) *Biotechniques*, 13, 188.
9. Rapley, R. (1994) *Mol. Biotechnol.* 2, 295–298.
10. Olszewski, M. et al. (2005) *Mol. Cell Probes*, 19, 203–205.
11. Baugh, L.R. et al. (2001) *Nucl. Acids Res.* 29, e29.
12. Villalva, C. et al. (2001) *Biotechniques* 31, 81–83, 86.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.