dATP Solution



1-800-632-7799 info@neb.com www.neb.com

N0440S

25 μmol Lot: 0531502 100 mM Store at -20°C Exo: 2/17

Description: Contains 0.25 ml of 100 mM ultrapure dATP.

Supplied in: Ultrapure water as a sodium salt at pH 7.4.

Concentration: The dATP Solution is supplied at a concentration of 100 mM (± 5 mM) as determined by A_{260} absorbance.

Diluent Compatibility: Can be diluted using sterile distilled water, preferably Milli-Q® water or can be diluted using sterile TE (10 mM Tris-HCl, 1 mM EDTA (pH 7.5).

Quality Controls

The purity of dATP is \geq 99% as determined by HPLC analysis.

0.5 kb, 2 kb and 5 kb Lambda PCR Assay – 25 cycles of PCR amplification of 1 ng Lambda DNA with 5 units of *Taq* DNA Polymerase in the presence of 200 μM dATP, dGTP, dCTP, and dTTP, 0.5 μM primers and 1X ThermoPol® Buffer results in the amplification of the specific 0.5 kb, 2 kb and 5 kb products as determined by agarose gel electrophoresis.

Phosphatase Activity Assay (pNPP Colorimetric Assay) – A protein phosphatase buffer solution containing 2 mM dATP Solution and 100 μ M p-nitrophenol phosphate, incubated for 4 hours at 37°C, yields no detectable phosphatase activity as determined by spectrophotometric analysis of released p-nitrophenylene anion at 405 nm.

Non-specific Nuclease Assay – A 50 μ l reaction in 1X NEBuffer 2 containing 1 μ g of T3 DNA or HindIII digested Lambda DNA and a minimum of 10 μ l of dATP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

For use with any DNA Polymerases from NEB.

Companion Products Sold Separately:

Deoxynucleotide Solution Mix

#N0447S 8 μ mol of each #0447L 40 μ mol of each

Deoxynucleotide Solution Set

#N0446S 25 µmol of each







NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

MILLI-Q® is a registered trademark of Millipore Corporation.

CERTIFICATE OF ANALYSIS

dATP Solution



1-800-632-7799 info@neb.com www.neb.com

N0440S

25 μmol Lot: 0531502 100 mM Store at -20°C Exp: 2/17

Description: Contains 0.25 ml of 100 mM ultrapure dATP.

Supplied in: Ultrapure water as a sodium salt at pH 7.4.

Concentration: The dATP Solution is supplied at a concentration of 100 mM (± 5 mM) as determined by A_{260} absorbance.

Diluent Compatibility: Can be diluted using sterile distilled water, preferably Milli-Q® water or can be diluted using sterile TE (10 mM Tris-HCl, 1 mM EDTA (pH 7.5).

Quality Controls

The purity of dATP is \geq 99% as determined by HPLC analysis.

0.5 kb, 2 kb and 5 kb Lambda PCR Assay – 25 cycles of PCR amplification of 1 ng Lambda DNA with 5 units of *Taq* DNA Polymerase in the presence of 200 μM dATP, dGTP, dCTP, and dTTP, 0.5 μM primers and 1X ThermoPol® Buffer results in the amplification of the specific 0.5 kb, 2 kb and 5 kb products as determined by agarose gel electrophoresis.

Phosphatase Activity Assay (pNPP Colorimetric Assay) – A protein phosphatase buffer solution containing 2 mM dATP Solution and 100 μ M p-nitrophenol phosphate, incubated for 4 hours at 37°C, yields no detectable phosphatase activity as determined by spectrophotometric analysis of released p-nitrophenylene anion at 405 nm.

Non-specific Nuclease Assay - A 50 μ l reaction in 1X NEBuffer 2 containing 1 μ g of T3 DNA or HindIII digested Lambda DNA and a minimum of 10 μ l of dATP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

For use with any DNA Polymerases from NEB.

Companion Products Sold Separately:

Deoxynucleotide Solution Mix

#N0447S 8 μ mol of each #N0447L 40 μ mol of each

Deoxynucleotide Solution Set

#N0446S 25 µmol of each







NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

MILLI-Q® is a registered trademark of Millipore Corporation.