

Lambda DNA– BstEII Digest



1-800-632-7799
info@neb.com
www.neb.com



N3014S 081120414041

N3014S

150 gel lanes (150 µg) Lot: 0811204 Exp: 4/14

500 µg/ml Store at –20°C

1.5 ml Gel Loading

Dye, Blue (6X) Store at 25°C

Description: The BstEII digest of lambda DNA (*cl857 ind 1 Sam 7*) yields 14 fragments suitable for use as molecular weight standards for agarose gel electrophoresis (1)

Supplied in: 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA.

Source: The phage is isolated from the heat-inducible lysogen *E. coli* λ *cl857 S7* and then isolated from the purified phage by phenol extraction and dialyzed. The double-stranded DNA is digested to completion with BstEII, phenol extracted and dialyzed against 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA.

Reagents supplied:
6X Gel Loading Dye, Blue


1X Gel Loading Dye, Blue:
2.5% Ficoll-400
11 mM EDTA
3.3 mM Tris-HCl (pH 8.0@25°C)
0.017% SDS
0.015% bromophenol blue

Usage Recommendation: The approximate mass of DNA in each of the bands in our Lambda DNA-BstEII Digest is as follows (assuming a 1.0 µg loading):

Fragment	Base Pairs	DNA Mass
1	8,454	174 ng
2	7,242	149 ng
3	6,369	131 ng
4	5,686	117 ng
5	4,822	99 ng
6	4,324	89 ng
7	3,675	76 ng
8	2,323	48 ng
9	1,929	40 ng
10	1,371	28 ng
11	1,264	26 ng
12	702	14 ng
13	224	5 ng
14	117	2 ng

Note: For long term storage, store at –20°C. If samples need to be diluted, use TE or other buffer of minimal ionic strength. DNA may denature if diluted in dH₂O and subsequently heated. Temperatures > 60°C may cause denaturation.

The cohesive ends of fragments 1 and 4 may be separated by heating to 60°C for 3 minutes.



(see other side)

CERTIFICATE OF ANALYSIS

Lambda DNA– BstEII Digest



1-800-632-7799
info@neb.com
www.neb.com



N3014S 081120414041

N3014S

150 gel lanes (150 µg) Lot: 0811204 Exp: 4/14

500 µg/ml Store at –20°C

1.5 ml Gel Loading

Dye, Blue (6X) Store at 25°C

Description: The BstEII digest of lambda DNA (*cl857 ind 1 Sam 7*) yields 14 fragments suitable for use as molecular weight standards for agarose gel electrophoresis (1)

Supplied in: 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA.

Source: The phage is isolated from the heat-inducible lysogen *E. coli* λ *cl857 S7* and then isolated from the purified phage by phenol extraction and dialyzed. The double-stranded DNA is digested to completion with BstEII, phenol extracted and dialyzed against 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA.

Reagents supplied:
6X Gel Loading Dye, Blue


1X Gel Loading Dye, Blue:
2.5% Ficoll-400
11 mM EDTA
3.3 mM Tris-HCl (pH 8.0@25°C)
0.017% SDS
0.015% bromophenol blue

Usage Recommendation: The approximate mass of DNA in each of the bands in our Lambda DNA-BstEII Digest is as follows (assuming a 1.0 µg loading):

Fragment	Base Pairs	DNA Mass
1	8,454	174 ng
2	7,242	149 ng
3	6,369	131 ng
4	5,686	117 ng
5	4,822	99 ng
6	4,324	89 ng
7	3,675	76 ng
8	2,323	48 ng
9	1,929	40 ng
10	1,371	28 ng
11	1,264	26 ng
12	702	14 ng
13	224	5 ng
14	117	2 ng

Note: For long term storage, store at –20°C. If samples need to be diluted, use TE or other buffer of minimal ionic strength. DNA may denature if diluted in dH₂O and subsequently heated. Temperatures > 60°C may cause denaturation.

The cohesive ends of fragments 1 and 4 may be separated by heating to 60°C for 3 minutes.



(see other side)

CERTIFICATE OF ANALYSIS

Suggested protocol for loading a sample:

The following protocol is recommended for a 5 mm wide lane.

1. Prepare loading mixture:

Distilled water	3 μ l
6X Blue Loading Dye	1 μ l
DNA Ladder	<u>2 μl</u>
Total volume	6 μ l

2. Mix gently

3. Load onto the agarose gel

Note: The components of the mixture should be scaled up or down, depending on the width of the agarose gel.

References:

1. Daniels, D.L. et al. (1983). In R.W. Hendrix, J.W. Roberts, F.W. Stahl and R.A. Weisberg (Eds.), *Lambda-II* (pp. 519–676). New York: Cold Spring Harbor Laboratory Press.
2. Forster, A.C. et al. (1985) *Nucl. Acids Res.* 13, 745–761.

Suggested protocol for loading a sample:

The following protocol is recommended for a 5 mm wide lane.

1. Prepare loading mixture:

Distilled water	3 μ l
6X Blue Loading Dye	1 μ l
DNA Ladder	<u>2 μl</u>
Total volume	6 μ l

2. Mix gently

3. Load onto the agarose gel

Note: The components of the mixture should be scaled up or down, depending on the width of the agarose gel.

References:

1. Daniels, D.L. et al. (1983). In R.W. Hendrix, J.W. Roberts, F.W. Stahl and R.A. Weisberg (Eds.), *Lambda-II* (pp. 519–676). New York: Cold Spring Harbor Laboratory Press.
2. Forster, A.C. et al. (1985) *Nucl. Acids Res.* 13, 745–761.