

PNGase F, Recombinant



1-800-632-7799
info@neb.com
www.neb.com



P0708S 002160118011

P0708S




15,000 units Lot: 0021601 Exp: 1/18

500,000 U/ml Store at **-20°C**

Description: Peptide: N-Glycosidase F, also known as PNGase F, is a recombinant amidase which cleaves between the innermost GlcNAc and asparagine residues of high mannose, hybrid, and complex oligosaccharides from N-linked glycoproteins (1).

Specificity:



Source: Cloned from *Elizabethkingia miricola* (formerly *Flavobacterium meningosepticum*) and expressed in *E. coli* (2).

Applications:

- Removal of carbohydrate residues from proteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C), 5 mM Na₂EDTA and 50% glycerol.

Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer:
(5% SDS, 0.4 M DTT)

10X GlycoBuffer 2:
[0.5 M Sodium Phosphate (pH 7.5 @ 25°C)]

10% NP-40

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

Reaction Conditions:

Typical reaction conditions are as follows:

- Combine 1–20 µg of glycoprotein, 1 µl of 10X Glycoprotein Denaturing Buffer and H₂O (if necessary) to make a 10 µl total reaction volume.
- Denature glycoprotein by heating reaction at 100°C for 10 minutes.
- Make a total reaction volume of 20 µl by adding 2 µl 10X GlycoBuffer 2, 2 µl 10% NP-40, H₂O and 1–2 µl PNGase F, Recombinant.
- Incubate reaction at 37°C for 1 hour.

Note: Reactions may be scaled-up linearly to accommodate larger reaction volumes.

MolecularWeight: 36,000 daltons.

Heat Inactivation: 500 units of enzyme were inactivated by incubation at 75°C for 10 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl. (65 NEB units = 1 IUB milliunit).

Unit Definition Assay: 10 µg of RNase B are denatured with 1X Glycoprotein Denaturing Buffer at 100°C for 10 minutes. After the addition of NP-40 and GlycoBuffer 2, two-fold dilutions of PNGase F, Recombinant are added and the reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized by SDS-PAGE.

Quality Assurance: No contaminating exoglycosidase or endoglycosidase activity could be detected. No contaminating proteolytic activity could be detected.

(see other side)

CERTIFICATE OF ANALYSIS

PNGase F, Recombinant



1-800-632-7799
info@neb.com
www.neb.com



P0708S 002160118011

P0708S




15,000 units Lot: 0021601 Exp: 1/18

500,000 U/ml Store at **-20°C**

Description: Peptide: N-Glycosidase F, also known as PNGase F, is a recombinant amidase which cleaves between the innermost GlcNAc and asparagine residues of high mannose, hybrid, and complex oligosaccharides from N-linked glycoproteins (1).

Specificity:



Source: Cloned from *Elizabethkingia miricola* (formerly *Flavobacterium meningosepticum*) and expressed in *E. coli* (2).

Applications:

- Removal of carbohydrate residues from proteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C), 5 mM Na₂EDTA and 50% glycerol.

Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer:
(5% SDS, 0.4 M DTT)

10X GlycoBuffer 2:
[0.5 M Sodium Phosphate (pH 7.5 @ 25°C)]

10% NP-40

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

Reaction Conditions:

Typical reaction conditions are as follows:

- Combine 1–20 µg of glycoprotein, 1 µl of 10X Glycoprotein Denaturing Buffer and H₂O (if necessary) to make a 10 µl total reaction volume.
- Denature glycoprotein by heating reaction at 100°C for 10 minutes.
- Make a total reaction volume of 20 µl by adding 2 µl 10X GlycoBuffer 2, 2 µl 10% NP-40, H₂O and 1–2 µl PNGase F, Recombinant.
- Incubate reaction at 37°C for 1 hour.

Note: Reactions may be scaled-up linearly to accommodate larger reaction volumes.

MolecularWeight: 36,000 daltons.

Heat Inactivation: 500 units of enzyme were inactivated by incubation at 75°C for 10 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl. (65 NEB units = 1 IUB milliunit).

Unit Definition Assay: 10 µg of RNase B are denatured with 1X Glycoprotein Denaturing Buffer at 100°C for 10 minutes. After the addition of NP-40 and GlycoBuffer 2, two-fold dilutions of PNGase F, Recombinant are added and the reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized by SDS-PAGE.

Quality Assurance: No contaminating exoglycosidase or endoglycosidase activity could be detected. No contaminating proteolytic activity could be detected.

(see other side)

CERTIFICATE OF ANALYSIS

Quality Controls

Glycosidase Assays: 5,000 units of PNGase F, Recombinant were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 µl reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

No other glycosidase activities were detected (ND) with the following substrates:

β-N-Acetylglucosaminidase:

GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC ND

β-N-Acetylgalactosaminidase:

GalNAcβ1-4Galβ1-4Glc-AMC ND

α-N-Acetylgalactosaminidase:

GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC ND

α-Fucosidase:

Galβ1-4 (Fucα1-3)GlcNAcβ1-3Galβ1-4Glc-AMC ND

Fucα1-2Galβ1-4Glc-AMC ND

Quality Controls

Glycosidase Assays: 5,000 units of PNGase F, Recombinant were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 µl reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

No other glycosidase activities were detected (ND) with the following substrates:

β-N-Acetylglucosaminidase:

GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC ND

β-N-Acetylgalactosaminidase:

GalNAcβ1-4Galβ1-4Glc-AMC ND

α-N-Acetylgalactosaminidase:

GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC ND

α-Fucosidase:

Galβ1-4 (Fucα1-3)GlcNAcβ1-3Galβ1-4Glc-AMC ND

Fucα1-2Galβ1-4Glc-AMC ND

β-Galactosidase:

Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC ND

Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC ND

α-Galactosidase:

Galα1-3Galβ1-4Gal-AMC ND

Galα1-6Galα1-6Glcα1-2Fru-AMC ND

α-Neuraminidase:

Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC ND

α-Mannosidase:

Manα1-3Manβ1-4GlcNAc-AMC ND

Manα1-6Manα1-6(Manα1-3)Man-AMC ND

α-Glucosidase: Glcα1-6Glcα1-4Glc-AMC ND

β-Xylosidase:

Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC ND

β-Mannosidase:

Manβ1-4Manβ1-4Man-AMC ND

Endo F₁, F₂, H:

Dansylated invertase high mannose. ND

β-Galactosidase:

Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC ND

Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC ND

α-Galactosidase:

Galα1-3Galβ1-4Gal-AMC ND

Galα1-6Galα1-6Glcα1-2Fru-AMC ND

α-Neuraminidase:

Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC ND

α-Mannosidase:

Manα1-3Manβ1-4GlcNAc-AMC ND

Manα1-6Manα1-6(Manα1-3)Man-AMC ND

α-Glucosidase: Glcα1-6Glcα1-4Glc-AMC ND

β-Xylosidase:

Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC ND

β-Mannosidase:

Manβ1-4Manβ1-4Man-AMC ND

Endo F₁, F₂, H:

Dansylated invertase high mannose. ND

Endo F₂, F₃:

Dansylated fibrinogen biantennary. ND

Protease Assay: After incubation of 10,000 units of PNGase F, Recombinant with 0.2 nmol of a standardized mixture of proteins in a 20 µl reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Notes: To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Since PNGase F, Recombinant activity is inhibited by SDS, it is essential to have NP-40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time.

PNGase F, Recombinant will not cleave *M*-linked glycans containing core α1-3 Fucose.

Endo F₂, F₃:

Dansylated fibrinogen biantennary. ND

Protease Assay: After incubation of 10,000 units of PNGase F, Recombinant with 0.2 nmol of a standardized mixture of proteins in a 20 µl reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Notes: To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Since PNGase F, Recombinant activity is inhibited by SDS, it is essential to have NP-40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time.

PNGase F, Recombinant will not cleave *M*-linked glycans containing core α1-3 Fucose.

References:

1. Maley, F. et al. (1989) *Anal. Biochem.* 180, 195–204.
2. Chen, M. New England Biolabs, Inc., unpublished results.

Companion Products:

RNase B

#P7817S 250 µg

Endoglycosidase Reaction Buffer Pack

B0701S 4 x 1 ml



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

References:

1. Maley, F. et al. (1989) *Anal. Biochem.* 180, 195–204.
2. Chen, M. New England Biolabs, Inc., unpublished results.

Companion Products:

RNase B

#P7817S 250 µg

Endoglycosidase Reaction Buffer Pack

B0701S 4 x 1 ml



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.