

E. coli Ribosome



1-800-632-7799
info@neb.com
www.neb.com



P0763S 007160618061

P0763S

1 mg **13.3 μM** **Lot: 0071606**

RECOMBINANT **Store at -80°C** **Exp: 6/18**

Description: The ribosome is the machinery for protein synthesis in a cell. The 70S ribosome consists of a small subunit (30S) and a large subunit (50S). This preparation of ribosome is highly active in the PURExpress® Protein Synthesis Kit (NEB #E6800).

E. coli Ribosome



1-800-632-7799
info@neb.com
www.neb.com



P0763S 007160618061

P0763S

1 mg **13.3 μM** **Lot: 0071606**

RECOMBINANT **Store at -80°C** **Exp: 6/18**

Description: The ribosome is the machinery for protein synthesis in a cell. The 70S ribosome consists of a small subunit (30S) and a large subunit (50S). This preparation of ribosome is highly active in the PURExpress® Protein Synthesis Kit (NEB #E6800).

Supplied in: 20 mM HEPES-KOH (pH 7.6), 10 mM MgAC₂, 30 mM KCl, and 7 mM β-mercaptoethanol.

Concentration:
33.3 mg/ml (13.3 μM)
1 A₂₆₀ = 60 μg/ml
1 mg/ml = 0.4 μM

Source: The ribosome is purified from an *E. coli* B strain.

Applications:

- *In vitro* translation
- Ribosome structural and functional studies
- Target for drug screening
- Starting material for isolation of native ribosomal RNAs (5S, 16S and 23S)

Quality Control Assays

E. coli Ribosome is functionally tested in the PURExpress Protein Synthesis Kit (NEB #E6800).

Supplied in: 20 mM HEPES-KOH (pH 7.6), 10 mM MgAC₂, 30 mM KCl, and 7 mM β-mercaptoethanol.

Concentration:
33.3 mg/ml (13.3 μM)
1 A₂₆₀ = 60 μg/ml
1 mg/ml = 0.4 μM

Source: The ribosome is purified from an *E. coli* B strain.

Applications:

- *In vitro* translation
- Ribosome structural and functional studies
- Target for drug screening
- Starting material for isolation of native ribosomal RNAs (5S, 16S and 23S)

Quality Control Assays

E. coli Ribosome is functionally tested in the PURExpress Protein Synthesis Kit (NEB #E6800).

Usage Notes: Avoid multiple freeze/thaw cycles. If you plan to use the ribosome multiple times, we recommend preparing aliquots after first thaw. Handle the ribosome gently, mix by tapping the tube, and avoid vigorous vortexing or high speed centrifugation. Mg²⁺ is necessary to keep the ribosome intact. If you are going to dilute the ribosome, be sure to include 10 mM MgAC₂ in the buffer.

Companion Products:

PURExpress *In Vitro* Protein Synthesis Kit
#E6800S 10 rxns
#E6800L 100 rxns

PURExpress Δ Ribosome Kit
#E3313S 10 rxns

PURExpress Δ RF123 Kit
#E6850S 10 rxns

PURExpress Disulfide Bond Enhancer
#E6820S 50 rxns

PURExpress Δ (aa, tRNA) Kit
#E6840S 10 rxns

Usage Notes: Avoid multiple freeze/thaw cycles. If you plan to use the ribosome multiple times, we recommend preparing aliquots after first thaw. Handle the ribosome gently, mix by tapping the tube, and avoid vigorous vortexing or high speed centrifugation. Mg²⁺ is necessary to keep the ribosome intact. If you are going to dilute the ribosome, be sure to include 10 mM MgAC₂ in the buffer.

Companion Products:

PURExpress *In Vitro* Protein Synthesis Kit
#E6800S 10 rxns
#E6800L 100 rxns

PURExpress Δ Ribosome Kit
#E3313S 10 rxns

PURExpress Δ RF123 Kit
#E6850S 10 rxns

PURExpress Disulfide Bond Enhancer
#E6820S 50 rxns

PURExpress Δ (aa, tRNA) Kit
#E6840S 10 rxns



NEW ENGLAND BIOLABS® and PUREXPRESS® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® and PUREXPRESS® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS