

Trypsin-ultra™, Mass Spectrometry Grade



1-800-632-7799
info@neb.com
www.neb.com



P8101S 007150916091

P8101S

100 µg Lot: 0071509 Exp: 9/16

5 x 20 µg Store at -20°C

Description: Trypsin-ultra, Mass Spectrometry Grade is a serine endopeptidase. It selectively cleaves peptide bonds C-terminal to lysine and arginine residues (1). Trypsin-ultra is treated with L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK) to inactivate any remaining chymotryptic activity. It is modified by acetylation of the ε-amino groups of lysine residues to prevent autolysis. Trypsin-ultra cleaves at Lys-Pro and Arg-Pro bonds at a much slower rate than other amino acid residues (2).

Source: Isolated from bovine (*Bos taurus*) pancreas

Applications:

- Digestion of proteins for proteomic analysis by Mass Spectrometry
- Protein and peptide identification

Reaction Conditions: 1X Trypsin-ultra, Reaction Buffer. Incubate at 37°C.

Reagents Supplied with Enzyme:

2X Trypsin-ultra, Reaction Buffer.

1X Trypsin-ultra, Reaction Buffer:

50 mM Tris-HCl
20 mM CaCl₂
pH 8.0 @ 25°C

Note: Substrate must be in phosphate-free buffer to prevent calcium precipitation with both reconstituted enzyme and enzyme buffer.

Molecular Weight: 23,675 daltons

Reconstitution: Trypsin-ultra, Mass Spectrometry Grade should be reconstituted by the addition of 20–200 µl of high purity water. Rapid autolysis is a function of enzyme concentration.

Storage Conditions: Supplied freeze-dried from a sodium acetate and calcium chloride buffer. Store at -20°C.

Can be stored frozen in solution at -20°C for up to 2 weeks. A decrease in activity will occur if stored in solution. Use only freshly reconstituted protease for best results.

Quality Assurance: Trypsin-ultra, Mass Spectrometry Grade is free of glycerol and detergents which may interfere with Matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) Mass Spectrometry (MS) or liquid chromatography (LC) methods.

Quality Controls

Functional Test (Calcitonin Peptide Digestion): A 20 µl reaction in Trypsin-ultra, Mass Spectrometry Grade Reaction Buffer containing 2 µg of human calcitonin peptide and 0.1 µg of Trypsin incubated for 16 hours at 37°C results in the expected digestion products determined by MALDI-TOF MS analysis.

Functional Test (Cytochrome C Digestion): A 20 µl reaction in Trypsin-ultra, Mass Spectrometry Grade Reaction Buffer containing 2 µg of Cytochrome C and 0.1 µg of Trypsin incubated for 16 hours at 37°C results in the expected digestion products determined by MALDI-TOF MS analysis.

Specific Activity: The specific activity of Trypsin-ultra, Mass Spectrometry Grade is between 1.8 µmol min⁻¹ mg⁻¹ and 3.0 µmol min⁻¹ mg⁻¹.

Note: Trypsin-ultra, Mass Spectrometry Grade is acetylated on multiple lysine residues. This protein appears as a single band on SDS-PAGE. This sequence is also available at www.neb.com.

(see other side)

CERTIFICATE OF ANALYSIS

Trypsin-ultra™, Mass Spectrometry Grade



1-800-632-7799
info@neb.com
www.neb.com



P8101S 007150916091

P8101S

100 µg Lot: 0071509 Exp: 9/16

5 x 20 µg Store at -20°C

Description: Trypsin-ultra, Mass Spectrometry Grade is a serine endopeptidase. It selectively cleaves peptide bonds C-terminal to lysine and arginine residues (1). Trypsin-ultra is treated with L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK) to inactivate any remaining chymotryptic activity. It is modified by acetylation of the ε-amino groups of lysine residues to prevent autolysis. Trypsin-ultra cleaves at Lys-Pro and Arg-Pro bonds at a much slower rate than other amino acid residues (2).

Source: Isolated from bovine (*Bos taurus*) pancreas

Applications:

- Digestion of proteins for proteomic analysis by Mass Spectrometry
- Protein and peptide identification

Reaction Conditions: 1X Trypsin-ultra, Reaction Buffer. Incubate at 37°C.

Reagents Supplied with Enzyme:

2X Trypsin-ultra, Reaction Buffer.

1X Trypsin-ultra, Reaction Buffer:

50 mM Tris-HCl
20 mM CaCl₂
pH 8.0 @ 25°C

Note: Substrate must be in phosphate-free buffer to prevent calcium precipitation with both reconstituted enzyme and enzyme buffer.

Molecular Weight: 23,675 daltons

Reconstitution: Trypsin-ultra, Mass Spectrometry Grade should be reconstituted by the addition of 20–200 µl of high purity water. Rapid autolysis is a function of enzyme concentration.

Storage Conditions: Supplied freeze-dried from a sodium acetate and calcium chloride buffer. Store at -20°C.

Can be stored frozen in solution at -20°C for up to 2 weeks. A decrease in activity will occur if stored in solution. Use only freshly reconstituted protease for best results.

Quality Assurance: Trypsin-ultra, Mass Spectrometry Grade is free of glycerol and detergents which may interfere with Matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) Mass Spectrometry (MS) or liquid chromatography (LC) methods.

Quality Controls

Functional Test (Calcitonin Peptide Digestion): A 20 µl reaction in Trypsin-ultra, Mass Spectrometry Grade Reaction Buffer containing 2 µg of human calcitonin peptide and 0.1 µg of Trypsin incubated for 16 hours at 37°C results in the expected digestion products determined by MALDI-TOF MS analysis.

Functional Test (Cytochrome C Digestion): A 20 µl reaction in Trypsin-ultra, Mass Spectrometry Grade Reaction Buffer containing 2 µg of Cytochrome C and 0.1 µg of Trypsin incubated for 16 hours at 37°C results in the expected digestion products determined by MALDI-TOF MS analysis.

Specific Activity: The specific activity of Trypsin-ultra, Mass Spectrometry Grade is between 1.8 µmol min⁻¹ mg⁻¹ and 3.0 µmol min⁻¹ mg⁻¹.

Note: Trypsin-ultra, Mass Spectrometry Grade is acetylated on multiple lysine residues. This protein appears as a single band on SDS-PAGE. This sequence is also available at www.neb.com.

(see other side)

CERTIFICATE OF ANALYSIS

Trypsin-ultra, Mass Spectrometry Grade Protein Sequence:

1 IVGGYTCAENSVPYQVSLNAGYHFCGGSLINDQWVVSAAHCYQYHIQVRLGEYNID
61 VLEGGEQFIDASKIIRHPKYSSWTLNDILLIKLSTPAVINARVSTLLLPSACASA
121 GTECLISGWGNTLSSGVNYPDLLQCLVAPLLSHADCEASYPGQITNNMICAGFLEG
181 GKDSCQGDSSGPPVACNGQLQGI VSWGYGCAQKKGKPGVYTKVCNYVDWIQETIAANS

References:

1. Northrop J. H. and Kunitz, M. (1931). *Science* 73, 262–263.
2. Perona J. J. and Craik, C.S. (1995). *Protein Sci.* 4, 337–360.




NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

TRYPSIN-ULTRA™ is a trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

MALDI-TOF MS: *Issatchenkia orientalis* Cytochrome c subjected to digestion by Trypsin-ultra, Mass Spectrometry Grade for 16 hours, dried and subjected to MALDI-TOF MS.



Trypsin-ultra, Mass Spectrometry Grade Protein Sequence:

1 IVGGYTCAENSVPYQVSLNAGYHFCGGSLINDQWVVSAAHCYQYHIQVRLGEYNID
61 VLEGGEQFIDASKIIRHPKYSSWTLNDILLIKLSTPAVINARVSTLLLPSACASA
121 GTECLISGWGNTLSSGVNYPDLLQCLVAPLLSHADCEASYPGQITNNMICAGFLEG
181 GKDSCQGDSSGPPVACNGQLQGI VSWGYGCAQKKGKPGVYTKVCNYVDWIQETIAANS

References:

1. Northrop J. H. and Kunitz, M. (1931). *Science* 73, 262–263.
2. Perona J. J. and Craik, C.S. (1995). *Protein Sci.* 4, 337–360.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

TRYPSIN-ULTRA™ is a trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

MALDI-TOF MS: *Issatchenkia orientalis* Cytochrome c subjected to digestion by Trypsin-ultra, Mass Spectrometry Grade for 16 hours, dried and subjected to MALDI-TOF MS.

