

New England Biolabs Product Specification

<i>Product Name:</i>	<i>hSMUG1</i>
<i>Catalog #:</i>	<i>M0336S/L</i>
<i>Concentration:</i>	<i>5,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to excise 1 pmol of deoxyuracil from a 34-mer oligonucleotide duplex containing a single dU site in a total reaction volume of 10 µl in 1 hour at 37°C.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton®X-100, 200 µg/ml BSA, (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0336S/L v1.0</i>
<i>Effective Date:</i>	<i>08 Jun 2018</i>

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of hSMUG1 incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 1 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of hSMUG1 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - hSMUG1 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



Date 08 Jun 2018

Derek Robinson
Director of Quality Control

