

New England Biolabs Product Specification

<i>Product Name:</i>	<i>BtsI-v2</i>
<i>Catalog #:</i>	<i>R0667S/L</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 55°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>50 mM KCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-R0667S/L v2.0</i>
<i>Effective Date:</i>	<i>21 May 2019</i>

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 50 units of BtsI-v2 incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of BtsI-v2 incubated for 15 minutes at 55°C results in complete digestion as determined by agarose gel electrophoresis.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda DNA with BtsI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with BtsI-v2.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 10 units of BtsI-v2 incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - BtsI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



Date 21 May 2019

Derek Robinson
Director of Quality Control

