

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	StickTogether™ DNA Ligase Buffer
Catalog Number:	B0535S
Concentration:	2 X Concentrate
Packaging Lot Number:	10114525
Expiration Date:	06/2023
Storage Temperature:	-20°C
Specification Version:	PS-B0535S v1.0
Composition (1X):	66 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, 1 mM ATP, 7.5 % PEG 6000, (pH 7.6 @ 25°C)

StickTogether™ DNA Ligase Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B0535AVIAL	StickTogether™ DNA Ligase Buffer	10077408	Pass	

Assay Name/Specification	Lot # 10114525
<b>pH (buffers/solutions)</b> The pH of 2X StickTogether™ DNA Ligase Buffer is between pH 7.5 and 7.7 at 25°C.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X StickTogether <sup>™</sup> DNA Ligase Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Functional Testing (DNA Ligase Buffer) A 20 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 50 ng of Lambda-HindIII DNA and 2,000 units of T4 DNA Ligase incubated for 5 minutes at 25°C results in approximately >95% ligation of the DNA fragments as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X StickTogether <sup>™</sup> DNA Ligase Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 40 ng of a 300	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10114525
base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90%	
of the substrate RNA remains intact as determined by fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 28 Jun 2021

Josh Hersey

Packaging Quality Control Inspector 28 Jun 2021

