

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: NEBuffer™ r2.1

Catalog Number: B6002S

Concentration: 10 X Concentrate

Packaging Lot Number: 10225734
Expiration Date: 06/2026
Storage Temperature: -20°C

Specification Version: PS-B6002S v1.0

Composition (1X): 10 mM Tris-HCl, 50 mM NaCl, 10 mM MgCl2, 100 µg/ml rAlbumin, (pH 7.9

@ 25°C)

NEBuffer™ r2.1 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B6002SVIAL	NEBuffer™ r2.1	10193045	Pass	

Assay Name/Specification	Lot # 10225734
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer™ r2.1 is between 55 and 62 mS/cm at 25°C.	
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 μl reaction in 1X NEBuffer™ r2.1 containing 1 μg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 μl reaction in 1X NEBuffer™ r2.1 containing 1 μg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



B6002S / Lot: 10225734

Page 1 of 2

Assay Name/Specification	Lot # 10225734
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer™ r2.1 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
pH (buffers/solutions) The pH of 10X NEBuffer™ r2.1 is between pH 7.8 and 8.0 at 25°C.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Nancy Considine Production Scientist 16 Jun 2023

Janey Gundan

Michael Tonello

Packaging Quality Control Inspector

26 Jan 2024



B6002S / Lot: 10225734

Page 2 of 2