

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: rCutSmart™ Buffer

Catalog Number: B6004S

Concentration: 10 X Concentrate

Packaging Lot Number: 10185541
Expiration Date: 03/2026
Storage Temperature: -20°C

Specification Version: PS-B6004S/V v1.0

Composition (1X): 50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium

Acetate, 100 μg/ml rAlbumin, (pH 7.9 @ 25°C)

rCutSmart™ Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B6004SVIAL	rCutSmart™ Buffer	10182170	Pass	

Assay Name/Specification	Lot # 10185541
Conductivity (buffers/solutions)	Pass
The conductivity of 10X rCutSmart™ Buffer is between 40 and 46 mS/cm at 25°C.	
Endonuclease Activity (Nicking, Buffer)	Pass
A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA	
incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and 1	
unit of Clal incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	
garage 2177 tab determined by agarese ger electropheresis.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of Mscl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA	
as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the	
substrate DNA as determined by agarose gel electrophoresis.	



B6004S / Lot: 10185541

Page 1 of 2

Assay Name/Specification	Lot # 10185541
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X rCutSmart™ Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
pH (buffers/solutions) The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Nancy Considine Production Scientist 06 Apr 2023

Janey Gundan

Josh Hersey Packaging Quality Control Inspector 02 May 2023



B6004S / Lot: 10185541