

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | rCutSmart™ Buffer |
|------------------------|---|
| Catalog Number: | B6004S |
| Concentration: | 10 X Concentrate |
| Packaging Lot Number: | 10197165 |
| Expiration Date: | 06/2026 |
| Storage Temperature: | -20°C |
| Specification Version: | PS-B6004S/V v1.0 |
| Composition (1X): | 50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 μg/ml rAlbumin, (pH 7.9 @ 25°C) |

| rCutSmart™ Buffer Component List | | | | |
|----------------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| B6004SVIAL | rCutSmart™ Buffer | 10196036 | Pass | |

| Assay Name/Specification | Lot # 10197165 |
|---|----------------|
| Conductivity (buffers/solutions) The conductivity of 10X rCutSmart [™] Buffer is between 40 and 46 mS/cm at 25°C. | Pass |
| Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X rCutSmart [™] Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart [™] Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart [™] Buffer containing 1 µg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart [™] Buffer containing 1 µg of Lambda dam- DNA and 1 unit of Clal incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10197165 |
|---|----------------|
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X rCutSmart [™] Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| RNase Activity (Buffer) A 10 µl reaction in 1X rCutSmart [™] Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |
| pH (buffers/solutions) The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C. | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

my Commun

Nancy Considine Production Scientist 16 Jun 2023

Michae 11.

Michael Tonello Packaging Quality Control Inspector 31 Jul 2023

