

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: rCutSmart™ Buffer

Catalog Number: B6004S

Concentration: 10 X Concentrate

Packaging Lot Number: 10215583
Expiration Date: 10/2026
Storage Temperature: -20°C

Specification Version: PS-B6004S/V v1.0

Composition (1X): 50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium

Acetate, 100 μg/ml rAlbumin, (pH 7.9 @ 25°C)

rCutSmart™ Buffer Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
B6004SVIAL	rCutSmart™ Buffer	10207415	Pass	

Assay Name/Specification	Lot # 10215583
Conductivity (buffers/solutions)	Pass
The conductivity of 10X rCutSmart™ Buffer is between 40 and 46 mS/cm at 25°C.	
Endonuclease Activity (Nicking, Buffer) A 50 μl reaction in 1X rCutSmart™ Buffer containing 1 μg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and 1 unit of Clal incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 μl reaction in 1X rCutSmart™ Buffer containing 1 μg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 μl reaction in 1X rCutSmart™ Buffer containing 1 μg of Lambda DNA and 1 unit of Mscl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass



B6004S / Lot: 10215583

Page 1 of 2

**Pass** 

Assay Name/Specification	Lot # 10215583
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X rCutSmart™ Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Nancy Considine Production Scientist 24 Oct 2023

my Gondon

pH (buffers/solutions)

Michael Tonello

Packaging Quality Control Inspector

07 Nov 2023



B6004S / Lot: 10215583

Page 2 of 2