

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	rCutSmart™ Buffer
Catalog Number:	B6004S
Concentration:	10 X Concentrate
Packaging Lot Number:	10232267
Expiration Date:	02/2027
Storage Temperature:	-20°C
Specification Version:	PS-B6004S/V v1.0
Composition (1X):	50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 μg/ml rAlbumin, (pH 7.9 @ 25°C)

rCutSmart™ Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B6004SVIAL	rCutSmart™ Buffer	10224841	Pass	

Assay Name/Specification	Lot # 10232267
<b>Conductivity (buffers/solutions)</b> The conductivity of 10X rCutSmart <sup>™</sup> Buffer is between 40 and 46 mS/cm at 25°C.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X rCutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart <sup>™</sup> Buffer containing 1 µg of Lambda dam- DNA and 1 unit of Clal incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass





240 County Road Ipswich, MA 01938-2723 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10232267
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X rCutSmart <sup>™</sup> Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X rCutSmart™ Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>pH (buffers/solutions)</b> The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

any Commun

Nancy Considine Production Scientist 08 Feb 2024

Michae 711.

Michael Tonello Packaging Quality Control Inspector 12 Feb 2024

