240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Gel Loading Dye, Purple (6X)

 Catalog #:
 B7024S

 Concentration:
 6X Concentrate

 Lot #:
 0171801

 Assay Date:
 01/2018

 Expiration Date:
 01/2021

Storage Temp: 25°C

Composition (1X): 3.3 mM Tris-HCl, 10 mM EDTA, 2.5 % Ficoll® 400, 0.08 % SDS, 0.02 % Dye 1, 0.0008 % Dye 2, (pH 8.0 @ 25°C)

Specification Version: PS-B7024S v1.0 Effective Date: 16 Feb 2018

| Assay Name/Specification (minimum release criteria) | Lot #0171801 |
|---|--------------|
| Endonuclease Activity (Nicking) - A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 μl of Gel Loading Dye, Purple (6X) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 10 μl of Gel Loading Dye, Purple (6X) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart® Buffer containing 1 µl of digested 2-Log Ladder DNA and a minimum of 10 µl of Gel Loading Dye, Purple (6X) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Gel Loading Dye, Purple (6X) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |

Authorized by Derek Robinson 16 Feb 2018







Inspected by
Tony Spear-Alfonso
10 Jan 2018