

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Blue Protein Loading Dye

Catalog Number: B7703S
Packaging Lot Number: 10121371
Expiration Date: 09/2023
Storage Temperature: -20°C

Specification Version: PS-B7703S v2.0

Composition (1X): 187.5 mM Tris-HCl, 6 % (w/v) SDS, 30 % Glycerol, 0.03 % Bromophenol

Blue, (pH 6.8 @ 25°C)

| Blue Protein Loading Dye Component List |                          |            |                      |  |
|---|--------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                  | Component Description    | Lot Number | Individual QC Result |  |
| B7705SVIAL                              | 30X Reducing Agent       | 10081852   | Pass                 |  |
| B7703SVIAL                              | Blue Protein Loading Dye | 10093120   | Pass                 |  |

| Assay Name/Specification  | Lot # 10121371 |
|---|----------------|
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of 1 kb Plus DNA Ladder DNA and a minimum of 5 µl of Blue Protein Loading Dye incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.                  | Pass           |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Blue Protein Loading Dye is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass           |
| Electrophoretic Pattern The components of the Blue Protein Loading Dye are tested to ensure the banding pattern of an NEB protein ladder on a 10-20% Tris-Glycine gel shows discrete, clearly identifiable bands at each size fragment of the marker when stained with Coomassie Blue at a concentration of 0.1%.                       | Pass           |
| Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 μl of Blue Protein Loading Dye incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.  | Pass           |



B7703S / Lot: 10121371

Page 1 of 2



This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Michael Dalton

Production Scientist

17 Sep 2021

Michael Tonello

Packaging Quality Control Inspector

17 Sep 2021

B7703S / Lot: 10121371

Page 2 of 2