

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	S-adenosylmethionine (SAM)
Catalog #:	B9003S
Concentration:	32 mM
<i>Lot</i> #:	1211712
Assay Date:	12/2017
Expiration Date:	09/2018
Storage Temp:	-20°C
Composition (1X):	0.005 M Sulfuric Acid, 10 % Ethanol
Specification Version:	PS-B9003S v1.0
Effective Date:	09 Apr 2018

Assay Name/Specification (minimum release criteria)	
Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 5 μ l of S-adenosylmethionine (SAM) incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of PhiX174-HaeIII DNA and a minimum of 5 μ l of S-adenosylmethionine (SAM) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Restriction Digest (CpG Resistant, SAM) - A 20 μ l reaction in 1X NEBuffer 2 containing 1 μ g of Lambda DNA, 1 unit of M. SssI (CpG Methyltransferase), and 160 μ M S-adenosylmethionine (SAM) is incubated for 1 hour at 37°C. The resulting DNA is resistant to digestion with BstUI as determined by agarose gel electrophoresis.	Pass

otunen

Authorized by Derek Robinson 09 Apr 2018



Jean Bastove

Inspected by Jean Bastable 06 Dec 2017