

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: E.coli DNA Ligase

Catalog Number: M0205S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of 6 μg of Lambda-HindIII DNA in 30 minutes at 16°C in a

total reaction volume of 20 μl.

Lot Number: 10025798
Expiration Date: 11/2020
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 µg/ml BSA

, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0205S/L v2.0

E.coli DNA Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0205SVIAL	E.coli DNA Ligase	10028141	Pass	
B0205SVIAL	E. coli DNA Ligase Reaction Buffer	10029823	Pass	

Assay Name/Specification	Lot # 10025798
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and	Pass
a minimum of 50 units of E. coli DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of E. coli DNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 20 units of E. coli DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) E. coli DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie	Pass



M0205S / Lot: 10025798

Page 1 of 2

Assay Name/Specification	Lot # 10025798
Blue detection.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of E. coli DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of E. coli DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

an K brenzen

Mary Lorenzen
Production Scientist
15 Nov 2018

NEW ENGLAND BioLabs Inc.

Josh Hersey

Packaging Quality Control Inspector

06 Jun 2019

