

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Uracil-DNA Glycosylase (UDG) |
|------------------------|---|
| Catalog Number: | M0280L |
| Concentration: | 5,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³ H]-uracil in a 50 μl reaction containing 0.2 μg DNA (10 ⁴ -10 ⁵ cpm/μg) in 30 minutes at 37°C. |
| Packaging Lot Number: | 10097696 |
| Expiration Date: | 01/2023 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 100 μg/ml BSA |
| Specification Version: | PS-M0280S/L v1.0 |

| Uracil-DNA Glycosylase (UDG) Component List | | | | |
|---|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0280LVIAL | Uracil-DNA Glycosylase (UDG) | 10097697 | Pass | |
| B0280SVIAL | UDG Reaction Buffer | 10084484 | Pass | |

| Assay Name/Specification | Lot # 10097696 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10097696 |
|---|----------------|
| gel electrophoresis. | |
| Protein Purity Assay (SDS-PAGE) Uracil-DNA Glycosylase (UDG) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Diggins

Lauren Higgins Production Scientist 20 Jan 2021

The OU Michae

Michael Tonello Packaging Quality Control Inspector 20 Jan 2021

