

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: yDcpS Catalog Number: M0463S Concentration: 200,000 U/ml

Unit Definition: One unit is defined as the amount of yDcpS required to convert 50%

of a 500 nM m7G-capped substrate to a 5'-diphosphorylated form in a

total reaction volume of 20 µl in 1 hour at 37°C.

Packaging Lot Number: 10085511 Expiration Date: 05/2022 -20°C Storage Temperature:

10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 %Storage Conditions:

Glycerol, (pH 7.4 @ 25°C)

PS-M0463S v1.0 Specification Version:

yDcpS Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0463SVIAL	yDcpS	10085510	Pass	
B0463AVIAL	10X yDcpS Reaction Buffer	10044009	Pass	

Assay Name/Specification	Lot # 10085511
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in yDcpS Reaction Buffer containing 1 µg of a mixture of single and	Pass
double-stranded [3H] E. coli DNA and a minimum of 200 units of yDcpS incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Phosphatase Activity (pNPP)	Pass
A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 200 units of yDcpS incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	
spectrophotometric analysis.	
Protein Purity Assay (SDS-PAGE) yDcpS is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity Assay (4 Hour Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 200 units of yDcpS is incubated at 37°C. After incubation for 4	



M0463S / Lot: 10085511

Page 1 of 2

Assay Name/Specification	Lot # 10085511
hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
Endonuclease Activity (Nicking) A 50 µl reaction in yDcpS Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of yDcpS incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist

21 Oct 2020

Michael Tonello

Packaging Quality Control Inspector

21 Oct 2020



M0463S / Lot: 10085511

Page 2 of 2