

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	OneTaq® Hot Start 2X Master Mix with GC Buffer
Catalog Number:	M0485L
Concentration:	2 X Concentrate
Packaging Lot Number:	10068724
Expiration Date:	11/2021
Storage Temperature:	-20°C
Specification Version:	PS-M0485S/L v2.0
Composition (1X):	80 mM Tris-SO4 (pH 9.2 @ 25°C), 20 mM (NH4)2SO4, 2 mM MgSO4, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 5 % DMSO, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 25 units/ml OneTaq® Hot Start DNA Polymerase

OneTaq® Hot Start 2X Master Mix with GC Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0485SVIAL	OneTaq® Hot Start 2X Master Mix with GC Buffer	10065642	Pass	
B9026AVIAL	OneTaq® High GC Enhancer	10031487	Pass	

Assay Name/Specification	Lot # 10068724
Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) A 50 μl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 μM dNTPs including [³ H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction.	Pass
RNase Activity (Extended Digestion) A 10 μ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ I of OneTaq® Hot Start 2X Master Mix with GC Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
PCR Amplification (Hot Start 2 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10068724
PCR Amplification (Enhancer Dependent, >70% GC-rich, Master Mix) A 25 μ l reaction in 1X OneTaq® Hot Start Master Mix with GC Buffer and 20% OneTaq® High GC Enhancer in the presence of 0.2 μ M primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the enhancer-dependent production of the 627 bp product.	Pass
PCR Amplification (Buffer Dependent, >65% GC-rich, Master Mix) A 25 μ I reaction in 1X OneTaq® Hot Start Master Mix with GC Buffer and 0.2 μ M primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the buffer-dependent production of the 737 bp product.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X OneTaq® Hot Start Master Mix with GC Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

vastie Vazquez

Christie Vazquez Production Scientist 30 Jan 2020

Mich

Michael Tonello Packaging Quality Control Inspector 27 Mar 2020

