

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Q5® Hot Start High-Fidelity 2X Master Mix
Catalog Number:	M0494L
Concentration:	2 X Concentrate
Lot Number:	10032779
Expiration Date:	10/2020
Storage Temperature:	-20°C
Specification Version:	PS-M0494S/L v1.0
Composition (1X):	Proprietary

Q5® Hot Start High-Fidelity 2X Master Mix Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0494SVIAL	Q5® Hot Start High-Fidelity 2X Master Mix	10029429	Pass	

Assay Name/Specification	Lot # 10032779
<b>RNase Activity (Extended Digestion)</b> A 10 $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 $\mu$ l of Q5® Hot Start High-Fidelity 2X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>PCR Amplification (Hot Start, Human Genomic DNA, Master Mix)</b> A 25 μl reaction in 1X Q5® Hot Start High-Fidelity Master Mix and 0.5 μM primers containing 50 ng Human Genomic DNA for 25 cycles of PCR amplification results in the expected 665 bp product and a decrease in non-specific genomic bands after pre-incubation at room temperature for 1 hour, when compared to a non-hot start control reaction.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Q5® High-Fidelity DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 2 units of Q5® High-Fidelity DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10032779
contamination is ≤ 1 E. coli genome.	
Protein Purity Assay (SDS-PAGE) Q5® High-Fidelity DNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>PCR Amplification (20 kb Lambda DNA, Master Mix)</b> A 50 μl reaction in 1X Q5® Hot Start High-Fidelity Master Mix and 1.0 μM primers containing 10 ng Lambda DNA for 22 cycles of PCR amplification results in the expected 20 kb product.	Pass
<b>Endonuclease Activity (Nicking, Polymerase, dNTP)</b> A 50 µl reaction in NEBuffer 2 in the presence of 400 µM dNTPs containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of Q5® High-Fidelity DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 1X Q5® Hot Start High-Fidelity Master Mix containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>PCR Amplification (7 kb Human Genomic DNA, Master Mix)</b> A 50 μl reaction in 1X Q5® Hot Start High-Fidelity Master Mix and 0.5 μM primers containing 20 ng Human Genomic DNA for 30 cycles of PCR amplification results in the expected 7 kb product.	Pass

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso Production Scientist 21 Dec 2018

Michae 211

Michael Tonello Packaging Quality Control Inspector 11 Jan 2019

