

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Quick CIP
Catalog Number:	M0525L
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme that hydrolyzes 1 μ mol of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml in 1 minute at 37°C.
Packaging Lot Number:	10168539
Expiration Date:	10/2024
Storage Temperature:	-20°C
Storage Conditions:	25 mM Tris-HCl , 1 mM MgCl2 , 0.1 mM ZnCl2 , 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version:	PS-M0525S/L v1.0

Quick CIP Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0525LVIAL	Quick CIP	10161215	Pass	
B6004SVIAL	rCutSmart™ Buffer	10163560	Pass	

Assay Name/Specification	Lot # 10168539
RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Quick CIP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking) A 50 μ I reaction in CutSmart® Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum of 50 units of Quick CIP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10168539
A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of Quick CIP incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 18 Oct 2022

Michae

Michael Tonello Packaging Quality Control Inspector 18 Oct 2022

