

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Quick CIP |
|------------------------|---|
| Catalog Number: | M0525S |
| Concentration: | 5,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme that hydrolyzes 1 μmol of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml in 1 minute at 37°C. |
| Packaging Lot Number: | 10161214 |
| Expiration Date: | 02/2024 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 25 mM Tris-HCl , 1 mM MgCl2 , 0.1 mM ZnCl2 , 50 % Glycerol, (pH 7.5 @ 25°C) |
| Specification Version: | PS-M0525S/L v1.0 |

| Quick CIP Component List | | | | |
|--------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0525SVIAL | Quick CIP | 10139068 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10156430 | Pass | |

| Assay Name/Specification | Lot # 10161214 |
|---|----------------|
| RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Quick CIP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using gel electrophoresis using fluorescent detection. | Pass |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 μI reaction in NEBuffer 4 containing 1 μg of PhiX174-HaeIII DNA and a minimum | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10161214 |
|---|----------------|
| of 50 units of Quick CIP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel | |
| electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 31 Aug 2022

Erin Varney

Packaging Quality Control Inspector 31 Aug 2022

