

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Quick CIP
Catalog Number: M0525S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 μmol

of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml

in 1 minute at 37°C.

Packaging Lot Number: 10175798
Expiration Date: 10/2024
Storage Temperature: -20°C

Storage Conditions: 25 mM Tris-HCl , 1 mM MgCl2 , 0.1 mM ZnCl2 , 50 % Glycerol, (pH 7.5

@ 25°C)

Specification Version: PS-M0525S/L v1.0

| Quick CIP Component List | | | | |
|--------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0525SVIAL | Quick CIP | 10161213 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10173661 | Pass | |

| Assay Name/Specification | Lot # 10175798 |
|--|----------------|
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Quick CIP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using gel electrophoresis using fluorescent detection. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum of 50 units of Quick CIP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Endonuclease Activity (Nicking) | Pass |



M0525S / Lot: 10175798 Page 1 of 2

| Assay Name/Specification | Lot # 10175798 |
|--|----------------|
| A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 29 Dec 2022 Michael Tonello

Packaging Quality Control Inspector

29 Dec 2022



M0525S / Lot: 10175798

Page 2 of 2