

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EnGen® Cas9 NLS, S. pyogenes

Catalog Number:M0646MConcentration:20 μMUnit Definition:N/A

Packaging Lot Number: 10102327
Expiration Date: 08/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0646T/M v2.0

EnGen® Cas9 NLS, S. pyogenes Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0646MVIAL	EnGen® Cas9 NLS, S. pyogenes	10102328	Pass	
B6003SVIAL	NEBuffer™ r3.1	10110766	Pass	

Assay Name/Specification	Lot # 10102327
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Functional Testing (Targeted Digestion) A 20 µl reaction in NEBuffer 3.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM sgRNA, and 100 nM EnGen® Spy Cas9 NLS incubated for 1 hour at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and	Pass



M0646M / Lot: 10102327

Page 1 of 2

Assay Name/Specification	Lot # 10102327
double-stranded [³H] E. coli DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Protein Purity Assay (SDS-PAGE) EnGen® Spy Cas9 NLS is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist 17 Aug 2021 Michael Tonello

Packaging Quality Control Inspector

17 Aug 2021



M0646M / Lot: 10102327