

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	EnGen® Cas9 NLS, S. pyogenes
Catalog Number:	M0646T
Concentration:	20 μM
Unit Definition:	N/A
Packaging Lot Number:	10070592
Expiration Date:	11/2021
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-M0646T/M v2.0

EnGen® Cas9 NLS, S. pyogenes Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0646TVIAL	EnGen® Cas9 NLS, S. pyogenes	10058697	Pass	
B7203SVIAL	NEBuffer™ 3.1	10072160	Pass	

Assay Name/Specification	Lot # 10070592
Functional Testing (Targeted Digestion) A 20 µl reaction in NEBuffer 3.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM sgRNA, and 100 nM EnGen® Spy Cas9 NLS incubated for 1 hour at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) EnGen® Spy Cas9 NLS is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10070592 Pass
Endonuclease Activity (Nicking) A 50 μ I reaction in NEBuffer 3.1 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Bhairavi Jani Production Scientist 17 Jun 2020

m. M. Michae

Michael Tonello Packaging Quality Control Inspector 17 Jun 2020

