

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Vaccinia Capping System
Catalog #:	M2080B-MT
Concentration:	10,000 units/ml
Unit Definition:	One unit of Vaccinia Capping Enzyme is defined as the amount of enzyme required to incorporate 10 pmol of $(\alpha^{32}P)$ GTP into an 80 nt transcript in 1 hour at 37°C.
Lot #:	0351507
Assay Date:	07/2015
Expiration Date:	7/2017
Storage Temp:	-20°C
Storage Conditions:	100 mM NaCl , 20 mM Tris-HCl (pH 8.0), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 0.1 % Triton®X-100
Specification Version:	<i>PS-M2080S v1.0</i>
Effective Date:	13 Apr 2015

Assay Name/Specification (minimum release criteria)	Lot #0351507
Endonuclease Activity (Nicking) - A 50 µl reaction in Capping Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of Vaccinia Capping System incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in Capping Buffer containing 1 μ g of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 10 units of Vaccinia Capping System incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Protein Purity Assay (SDS-PAGE) - Vaccinia Capping System is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single- stranded RNA and a minimum of 10 units of Vaccinia Capping System is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

otim

Authorized by Derek Robinson 13 Apr 2015



Inspected by Bhairavi Jani 19 Aug 2015