

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Vaccinia Capping System

Catalog Number: M2080S
Concentration: 10,000 U/ml

Unit Definition: One unit of Vaccinia Capping Enzyme is defined as the amount of

enzyme required to incorporate 10 pmol of (α<sup>32</sup>P) GTP into an 80 nt

transcript in 1 hour at 37°C.

Lot Number: 10050813
Expiration Date: 02/2021
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 20 mM Tris-HCl (pH 8.0), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, 0.1 % Triton®X-100

Specification Version: PS-M2080S v1.0

Vaccinia Capping System Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
N2080AVIAL	GTP	10036923	Pass	
M2080SVIAL	Vaccinia Capping System	10036921	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10046083	Pass	
B2080AVIAL	10X Capping Buffer	10035769	Pass	

Assay Name/Specification	Lot # 10050813
RNase Activity (Extended Digestion) A 10 $\mu$ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Vaccinia Capping System is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Protein Purity Assay (SDS-PAGE)  Vaccinia Capping System is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in Capping Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 10 units of Vaccinia Capping System incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking)	Pass



M2080S / Lot: 10050813

Page 1 of 2

Assay Name/Specification	Lot # 10050813
A 50 µl reaction in Capping Buffer containing 1 µg of supercoiled PhiX174 DNA and a	
minimum of 10 units of Vaccinia Capping System incubated for 4 hours at 37°C results	
in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Bhairavi Jani

Production Scientist

30 Jan 2019

Michael Tonello

Packaging Quality Control Inspector

24 Jul 2019

