

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Vaccinia Capping System

Catalog Number: M2080S Concentration: 10,000 U/ml

Unit Definition: One unit of Vaccinia Capping Enzyme is defined as the amount of

enzyme required to incorporate 10 pmol of (α³²P) GTP into an 80 nt

transcript in 1 hour at 37°C.

Packaging Lot Number: 10217209 Expiration Date: 04/2025 Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl (pH 8.0), 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol , 0.1 % Triton®X-100

Specification Version: PS-M2080S v1.0

Vaccinia Capping System Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N2080AVIAL	GTP	10179511	Pass	
M2080SVIAL	Vaccinia Capping System	10182558	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10210241	Pass	
B2080AVIAL	10X Capping Buffer	10217210	Pass	

Assay Name/Specification	Lot # 10217209
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in Capping Buffer containing 1 µg of supercoiled PhiX174 DNA and a	
minimum of 10 units of Vaccinia Capping System incubated for 4 hours at 37°C results	
in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in Capping Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 10 units of Vaccinia Capping	
System incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Protein Purity Assay (SDS-PAGE)	Pass
Vaccinia Capping System is ≥ 95% pure as determined by SDS-PAGE analysis using	
Coomassie Blue detection.	
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	



M2080S / Lot: 10217209

Page 1 of 2

and a minimum of 400 miles of Manadala Constant On the minimum to the Late of a C700 Affine	Lot # 10217209
and a minimum of 10 units of Vaccinia Capping System is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jessica Cane Production Scientist 06 Apr 2023 Michael Tonello
Packaging Quality C

Packaging Quality Control Inspector



