

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	7-deaza-dGTP
Catalog Number:	N0445S
Concentration:	5 mM
Unit Definition:	N/A
Packaging Lot Number:	10232060
Expiration Date:	02/2026
Storage Temperature:	-20°C
Storage Conditions:	Supplied in Ultrapure water as a lithium salt , (pH 7.0)
Specification Version:	PS-N0445S/L v2.0

7-deaza-dGTP Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N0445SVIAL	7-deaza-dGTP	10226743	Pass	

Assay Name/Specification	Lot # 10232060
Endonuclease Activity (Nicking) A 50 μ I reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 20 μ I of 7-deaza-dGTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μ I reaction in NEBuffer 2 containing 1 μ g of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 μ I of 7-deaza-dGTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda DNA, 7-deaza) A 50 μ I reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μ M primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2 kb Lambda DNA, 7-deaza) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 2 kb product.	Pass





240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10232060
PCR Amplification (5 kb Lambda DNA, 7-deaza) A 50 μ I reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μ M primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
Phosphatase Activity (pNPP) A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 80 μl 7-deaza-dGTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Physical Purity (HPLC) 7-deaza-dGTP is \geq 95% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) A 10 μ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ I of 7-deaza-dGTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lea Antonopoulo

Lea Antonopoulos Production Scientist

Josh Hersey

Packaging Quality Control Inspector 14 Mar 2024

