

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: dUTP Solution

Catalog Number: N0459S
Concentration: 100 mM
Packaging Lot Number: 10187082
Expiration Date: 02/2025
Storage Temperature: -20°C

Storage Conditions: Supplied in Ultrapure water as a sodium salt (pH 7.5)

Specification Version: PS-N0459S v2.0

dUTP Solution Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N0459SVIAL	dUTP Solution	10180528	Pass	

Assay Name/Specification	Lot # 10187082
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 μl of dUTP Solution incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 2 containing 1 μg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 4 μl of dUTP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda, dNTPs) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dGTP, dCTP, and dUTP and 0.2 μM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2.0 kb Lambda, dNTPs) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dGTP, dCTP, and dUTP and 0.2 μM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.	Pass



N0459S / Lot: 10187082

Page 1 of 2

Assay Name/Specification	Lot # 10187082
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 16 µl of dUTP Solution incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Physical Purity (HPLC) dUTP Solution is ≥ 99% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of dUTP Solution is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 μl of dUTP Solution is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lea Antonpoulos

Production Scientist

03 Mar 2023

Michael Tonello

Packaging Quality Control Inspector

13 Jul 2023



N0459S / Lot: 10187082

Page 2 of 2