

New England Biolabs Certificate of Analysis

Product Name: α 1-2,3,6 Mannosidase
Catalog Number: P0768L
Concentration: 2,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the terminal mannose from 1 nmol of Man(α 1,3)-Man(β 1,4)-GlcNAc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Lot Number: 10008736
Expiration Date: 04/2019
Storage Temperature: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, (pH 7.5 @ 25°C)
Specification Version: PS-P0768S/L v1.0

α 1-2,3,6 Mannosidase Component List

NEB Part Number	Component Description	Lot Number	Individual QC Result
P0768LVIAL	α 1-2,3,6 Mannosidase	0111804	Pass
B1703SVIAL	10X Glycobuffer 4	0021710	Pass
B0768SVIAL	10X Zinc	0021804	Pass

Assay Name/Specification	Lot # 10008736
Glycosidase Activity (α-Neuraminidase) A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 4 units of α 1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-2 Fucosidase) A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-2Gal β 1-4Glc-AMC) and 4 unit of α 1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Fucosidase) A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 4 units of α 1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass

Assay Name/Specification	Lot # 10008736
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fuca1-2)Galβ1-4Glc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass

Assay Name/Specification	Lot # 10008736
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Glyco Buffer 4 containing 24 μg of a standard mixture of proteins and a minimum of 10 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Glycosidase Activity (Endo F1, F2, H) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (Endo F2, F3) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (PNGase F) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 μl reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 4 units of α1-2,3,6 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Alicia Bielik
Production Scientist
16 Aug 2018



Josh Hersey
Packaging Quality Control Inspector
16 Aug 2018