

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: HindIII

Catalog Number: R0104S

Concentration: 20,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10023264
Expiration Date: 06/2020
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0104S/L v1.0

HindIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0104SVIAL	HindIII	10012445	Pass	
B7202SVIAL	NEBuffer™ 2.1	0261805	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10021128	Pass	

Assay Name/Specification	Lot # 10023264
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of HindIII, religated	Pass
and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of HindIII incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [3H] E. coli DNA and a minimum of 200 units of HindIII incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 200-fold over-digestion of Lambda DNA with HindIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
Inaginents can be ligated with 14 DIVA ligase in 10 hours at 10°C. Of these ligated	



R0104S / Lot: 10023264

Page 1 of 2

Assay Name/Specification	Lot # 10023264
fragments, >95% can be recut with HindIII.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 60 Units of HindIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) HindIII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist

28 Jun 2018

Michael Tonello

Packaging Quality Control Inspector

05 Nov 2018



R0104S / Lot: 10023264

Page 2 of 2