

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: HaelII
Catalog Number: R0108T
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10149876
Expiration Date: 03/2024
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0108T/M v1.0

| HaellI Component List  |                              |            |                      |  |
|------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description        | Lot Number | Individual QC Result |  |
| R0108TVIAL             | HaellI                       | 10140449   | Pass                 |  |
| B7024AVIAL             | Gel Loading Dye, Purple (6X) | 10144740   | Pass                 |  |
| B6004SVIAL             | rCutSmart™ Buffer            | 10148729   | Pass                 |  |

| Assay Name/Specification  | Lot # 10149876 |
|---|----------------|
| Protein Purity Assay (SDS-PAGE)   | Pass           |
| HaelII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.  |                |
| Exonuclease Activity (Radioactivity Release)  | Pass           |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of HaeIII incubated for                                   |                |
| 4 hours at 37°C releases <0.1% of the total radioactivity.  |                |
| Ligation and Recutting (Terminal Integrity)   | Pass           |
| After a 20-fold over-digestion of Lambda DNA with HaeIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HaeIII. |                |
| Non-Specific DNase Activity (16 Hour)   | Pass           |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of HaeIII incubated for 16 hours at 37°C results in a DNA pattern free of                               |                |
| detectable nuclease degradation as determined by agarose gel electrophoresis.   |                |



R0108T / Lot: 10149876

Page 1 of 2



This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

18 May 2022

Erin Varney

Packaging Quality Control Inspector

18 May 2022



R0108T / Lot: 10149876

Page 2 of 2