

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Sfil

Catalog Number: R0123S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba in 1 hour at 50°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10093175
Expiration Date: 10/2022
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0123S/L v1.0

Sfil Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0123SVIAL	Sfil	10085829	Pass	
B7204SVIAL	CutSmart® Buffer	10091029	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089401	Pass	

Assay Name/Specification	Lot # 10093175
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Sfil incubated for 4 hours at 50°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of pXba DNA with Sfil, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Sfil.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 100 units of Sfil incubated for 16 hours at 50°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Sfil is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass



R0123S / Lot: 10093175

Page 1 of 2

Assay Name/Specification	Lot # 10093175
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 100 units of Sfil incubated for 4 hours at 50°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

13 Dec 2020

Josh Hersey

Packaging Quality Control Inspector

13 Dec 2020



R0123S / Lot: 10093175

Page 2 of 2