

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Hhal
Catalog Number:	R0139S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10218953
Expiration Date:	11/2025
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version:	PS-R0139S/L v2.0

Hhal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0139SVIAL	Hhal	10213831	Pass	
B6004SVIAL	rCutSmart™ Buffer	10209242	Pass	

Assay Name/Specification	Lot # 10218953
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 60 units of Hhal incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart [™] Buffer containing 1 µg of Lambda DNA and 1 µl of Hhal incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Hhal, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Hhal.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 60 units of Hhal incubated for 16 hours at 37°C results in a DNA pattern free of	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10218953
detectable nuclease degradation as determined by agarose gel electrophoresis.	
qPCR DNA Contamination (E. coli Genomic)	Pass
A minimum of 20 units of Hhal is screened for the presence of E. coli genomic DNA	
using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results	
are quantified using a standard curve generated from purified E. coli genomic DNA.	
The measured level of E. coli genomic DNA contamination is \leq 1 E. coli genome.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun

Production Scientist 27 Oct 2023

on ll Michae

Michael Tonello Packaging Quality Control Inspector 04 Dec 2023

