

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Smal
Catalog Number:	R0141L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (HindIII digest) in 1 hour at 25°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10087669
Expiration Date:	10/2022
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version:	PS-R0141S/L v1.0

Smal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0141LVIAL	Smal	10087670	Pass	
B7204SVIAL	CutSmart® Buffer	10089403	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089392	Pass	

Assay Name/Specification	Lot # 10087669
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of Smal, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of Smal incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Smal incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Smal, ~50% of the DNA fragments	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10087669
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Smal.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 60 Units of Smal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 03 Dec 2020

Josh Hersey

Packaging Quality Control Inspector 03 Dec 2020

