

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Bglll
Catalog Number: R0144S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number:10017161Expiration Date:03/2020Storage Temperature:-20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0144S/L v1.0

| BgIII Component List   |                              |            |                      |  |
|------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description        | Lot Number | Individual QC Result |  |
| R0144SVIAL             | BgIII                        | 0441803    | Pass                 |  |
| B7203SVIAL             | NEBuffer™ 3.1                | 10010189   | Pass                 |  |
| B7024SVIAL             | Gel Loading Dye, Purple (6X) | 10011445   | Pass                 |  |

| Assay Name/Specification   | Lot # 10017161 |
|--|----------------|
| Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of BgIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.                                     | Pass           |
| Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 Units of BglII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of BgIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.  | Pass           |
| Ligation and Recutting (Terminal Integrity)  After a 20-fold over-digestion of Lambda DNA with BgIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,   | Pass           |



R0144S / Lot: 10017161

Page 1 of 2

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso Production Scientist

06 Jun 2018

Michael Tonello

Packaging Quality Control Inspector

30 Jul 2018



R0144S / Lot: 10017161

Page 2 of 2