

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EcoRV
Catalog Number: R0195T
Concentration: 100,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10146778
Expiration Date: 03/2024
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0195T/M v1.0

| EcoRV Component List | | | |
|------------------------|------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0195TVIAL | EcoRV | 10146777 | Pass |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10132772 | Pass |
| B6003SVIAL | NEBuffer™ r3.1 | 10132774 | Pass |

| Assay Name/Specification | Lot # 10146778 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of EcoRV incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of EcoRV incubated for 4 hours at 37°C results in <50% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of EcoRV, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with EcoRV, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, | Pass |



R0195T / Lot: 10146778

Page 1 of 2

| Assay Name/Specification | Lot # 10146778 |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| >95% can be recut with EcoRV. | |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of Lambda DNA and a minimum of 20 units of EcoRV incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Protein Purity Assay (SDS-PAGE) EcoRV is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

06 Apr 2022

Michael Tonello

Packaging Quality Control Inspector

06 Apr 2022



R0195T / Lot: 10146778

Page 2 of 2