

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Mlul   |
|------------------------|--|
| Catalog Number:        | R0198S   |
| Concentration:         | 10,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Packaging Lot Number:  | 10066337   |
| Expiration Date:       | 11/2021  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R0198S/L v1.0   |

| Mlul Component List |                              |            |                      |  |
|---------------------|------------------------------|------------|----------------------|--|
| NEB Part Number     | Component Description        | Lot Number | Individual QC Result |  |
| R0198SVIAL          | Mlul                         | 10058399   | Pass                 |  |
| B7203SVIAL          | NEBuffer™ 3.1                | 10053974   | Pass                 |  |
| B7024SVIAL          | Gel Loading Dye, Purple (6X) | 10065745   | Pass                 |  |

| Assay Name/Specification  | Lot # 10066337 |
|---|----------------|
| <b>Endonuclease Activity (Nicking)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled pUC19 DNA and a<br>minimum of 30 units of Mlul incubated for 4 hours at 37°C results in <20% conversion<br>to the nicked form as determined by agarose gel electrophoresis.                 | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 100<br>Units of Mlul incubated for 16 hours at 37°C results in a DNA pattern free of<br>detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 10-fold over-digestion of Lambda DNA with Mlul, >95% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>>95% can be recut with Mlul.   | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Mlul incubated for 4  | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification                                 | Lot # 10066337 |
|--|----------------|
| hours at 37°C releases <0.1% of the total radioactivity. |                |

This product has been tested and shown to be in compliance with all specifications.

tephanie Onetto

Stephanie Cornelio Production Scientist 08 Nov 2019

Michae m. N.

Michael Tonello Packaging Quality Control Inspector 23 Mar 2020

