

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Rsrll
Catalog Number:	R0501S
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10230866
Expiration Date:	12/2024
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton® X-100, 200 μg/ml BSA (pH 7.4 @ 25°C)
Specification Version:	PS-R0501S/L v2.0

RsrII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0501SVIAL	Rsrll	10221057	Pass	
B6004SVIAL	rCutSmart™ Buffer	10228580	Pass	

Assay Name/Specification	Lot # 10230866
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of RsrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of RsrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with RsrII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with RsrII.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of Lambda DNA and a minimum of 50 units of RsrII incubated for 16 hours at 37⁰C results in a DNA pattern free of	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10230866
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE)	Pass
RsrII is $\geq$ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun

Production Scientist 19 Dec 2023

ton. ll Michae

Michael Tonello Packaging Quality Control Inspector 23 Feb 2024

