

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	ApaLl
Catalog Number:	R0507L
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (Hind III digest) in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10147391
Expiration Date:	04/2024
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0507S/L v1.0

ApaLI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0507LVIAL	ApaLI	10147392	Pass	
B6004SVIAL	rCutSmart™ Buffer	10143288	Pass	

Assay Name/Specification	Lot # 10147391
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled M13mp19 DNA and a minimum of 50 Units of ApaLI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of ApaLI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 Units of ApaLI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda-HindIII DNA with ApaLI, >95% of the DNA	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10147391
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with ApaLI.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 13 Apr 2022

Josh Hersey

Packaging Quality Control Inspector 13 Apr 2022

