

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | ApaLI  |
|------------------------|--|
| Catalog #:             | R0507S/L   |
| Concentration:         | 10,000 units/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA (Hind III digest) in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l. |
| Lot #:                 | 0401301  |
| Assay Date:            | 01/2013  |
| Expiration Date:       | 01/2015  |
| Storage Temp:          | -20 °C   |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA   |
| Specification Version: | PS-R0507S/L v1.0   |
| Effective Date:        | 10 Sep 2013  |

| Assay Name/Specification (minimum release criteria)  | Lot #0401301 |
|--|--------------|
| <b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled M13mp19 DNA and a minimum of 50 Units of ApaLI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.                              | Pass         |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of ApaLI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda-HindIII DNA with ApaLI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with ApaLI.   | Pass         |
| Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda-<br>HindIII DNA and a minimum of 100 Units of ApaLI incubated for 16 hours at 37°C results in a DNA pattern free<br>of detectable nuclease degradation as determined by agarose gel electrophoresis.        | Pass         |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

otunen

Authorized by Derek Robinson 10 Sep 2013



Diana doam

Inspected by Diana O'Loane 10 Sep 2013