

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Bsu36l
Catalog Number:	R0524L
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (Hind III digest) in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10160777
Expiration Date:	06/2024
Storage Temperature:	-20°C
Storage Conditions:	250 mM NaCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 0.15 % TritonX-100 , 200 μg/ml BSA
Specification Version:	PS-R0524S/L v1.0

Bsu36I Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0524LVIAL	Bsu36l	10155327	Pass	
B6004SVIAL	rCutSmart™ Buffer	10156434	Pass	

Assay Name/Specification	Lot # 10160777
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 30 units of Bsu36l incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Bsu36l incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of Bsu36l incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 2-fold over-digestion of Lambda-HindIII DNA with Bsu36I, ~25% of the DNA	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10160777
fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated	
fragments, >95% can be recut with Bsu36I.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 08 Sep 2022

Erin Varney

Packaging Quality Control Inspector 08 Sep 2022

