

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: BspEl
Catalog Number: R0540S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (dam) in 1 hour at 37°C in a total reaction volume of

50 μl.

Packaging Lot Number: 10090114
Expiration Date: 11/2022
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0540S/L v1.0

BspEl Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0540SVIAL	BspEl	10090112	Pass	
B7203SVIAL	NEBuffer™ 3.1	10085495	Pass	

Assay Name/Specification	Lot # 10090114
Blue-White Screening (Terminal Integrity) A sample of LITMUS38i vector linearized with a 10-fold excess of BspEI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene	Pass
results in <1% white colonies.  Endonuclease Activity (Nicking)  A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 Units of BspEl incubated for 4 hours at 37°C results in <20%	Pass
conversion to the nicked form as determined by agarose gel electrophoresis.  Exonuclease Activity (Radioactivity Release)	Pass
A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of BspEl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda dam- DNA with BspEI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass



R0540S / Lot: 10090114

Page 1 of 2

Assay Name/Specification	Lot # 10090114
fragments, >95% can be recut with BspEI.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda dam- DNA and a minimum of 50 units of BspEI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

09 Dec 2020

Michael Tonello

Packaging Quality Control Inspector

09 Dec 2020



R0540S / Lot: 10090114

Page 2 of 2