

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: DpnII
Catalog Number: R0543S
Concentration: 10,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of

50 μl.

Packaging Lot Number: 10154647
Expiration Date: 06/2024
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0543S/L v1.0

DpnII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0543SVIAL	DpnII	10154646	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10153339	Pass	
B0543SVIAL	NEBuffer™ DpnII	10146004	Pass	

Assay Name/Specification	Lot # 10154647
Protein Purity Assay (SDS-PAGE)	Pass
DpnII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer DpnII containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of DpnII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer DpnII containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of DpnII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda dam- DNA with DpnII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass



R0543S / Lot: 10154647 Page 1 of 2

Assay Name/Specification	Lot # 10154647
fragments, >95% can be recut with DpnII.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer DpnII containing 1 µg of Lambda dam- DNA and a minimum of 100 units of DpnII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

08 Jul 2022

Michael Tonello

Packaging Quality Control Inspector

08 Jul 2022

